

WHAT DO NETWORKS HAVE TO DO WITH CLIMATE?

BY ANASTASIOS A. TSONIS, KYLE L. SWANSON, AND PAUL J. ROEBBER

Advances in understanding coupling in complex networks offer new ways of studying the collective behavior of interactive systems and already have yielded new insights in many areas of science.

Last summer one of the coauthors (AAT) visited the Greek island of Corfu. During summer the population of the island is 100,000 people. Before he went there, he knew only two Corfians. The first night of his stay he dined in one of the restaurants close to his hotel. Greeks, being the friendly people they are, always open a discussion about who you are, what you do, and so on. As the waiter took his order they started talking about such things. What do you think the probability is that the waiter knew one of the two above-mentioned Corfians? This problem has an analytical solution, but the fact is that the waiter did know one of the two Corfians. Otherwise stated, it only took one connection between two people that

did not know each other to arrive at a common link. It is a small world after all! Through the work of the American psychologist Stanley Milgram and other subsequent investigations we know that any two of the six billion people on Earth are linked by a trail of only six people (Milgram 1967). This is referred to as the *six degrees of separation*.

Insights in such strange, but otherwise common, connections have been provided by the study of networks. A network is a system of interacting agents. In the literature an agent is called a node. The nodes in a network can be anything. For example, in the network of actors, the nodes are actors that are connected to other actors with whom they have appeared in a movie. In a network of species the nodes are species that are connected to other species with which they interact. In the network of scientists, the nodes are scientists that are connected to other scientists with whom they have collaborated. In the grand network of humans each node is an individual who is connected to people he or she knows.

There are four basic types of networks, which are described below.

Regular (ordered) networks. These networks have a fixed number of nodes, with each node having the same number of links connecting it in a specific way to a number of neighboring nodes (Fig. 1, left

AFFILIATIONS: TSONIS, SWANSON, AND ROEBBER—Department of Mathematical Sciences, Atmospheric Sciences Group, University of Wisconsin—Milwaukee, Milwaukee, Wisconsin

CORRESPONDING AUTHOR: Anastasios Tsonis, Department of Mathematical Sciences, Atmospheric Sciences Group, University of Wisconsin—Milwaukee, Milwaukee, WI 53201-0413
E-mail: aatsonis@uwm.edu

The abstract for this article can be found in this issue, following the table of contents.

DOI:10.1175/BAMS-87-5-585

In final form 10 January 2006
©2006 American Meteorological Society

panel). If each node is linked to all other nodes in the network, then the network is a fully connected network.

Classical random networks. In these networks (Erdos and Renyi 1960) the nodes are connected at random (Fig. 1, right panel). In this case the degree distribution is a Poisson distribution (the degree distribution p_k gives the probability that a node in the network is connected to k other nodes). The problem with these networks is that they are not very stable. Removal of a number of nodes at random may fracture the network to noncommunicating parts.

Small-world networks. Regular networks are locally clustered, which means that, unless they are fully wired, it takes many steps to go from one node to another node away from its immediate neighborhood. On the contrary, random networks do not exhibit local clustering. Faraway nodes can be connected as easily as nearby nodes. In this case, information may be transported all over the network much more efficiently than in ordered networks. Thus, random networks exhibit efficient information transfer and regular networks do not (unless they are fully connected). This dichotomy of networks as being either regular or random is undesirable, because one could expect that in nature networks should be efficient in processing information and at the same time be stable. Work in this direction led to a new type of network, which was proposed a few years ago by the American mathematicians Duncan Watts and Steven Strogatz (1998), that is called a “small world” network. A small-world network is a superposition of regular and classical random graphs. Such networks exhibit a high degree of local clustering, but a small number of long-range connections makes them as efficient in transferring information as random networks. Those long-range connections do not have to be designed. A few long-range connections added at random will do the trick (Fig. 1, middle panel). Both random and small-world networks are rather homogeneous networks in which each node has approximately the same number of links $\langle k \rangle$. Both have nearby Poisson degree distributions that peak at $\langle k \rangle$ and decay exponentially for large k .

Networks with a given degree distribution. The small-world architecture can explain phenomena such as




FIG. 1. Illustration of a regular, a small-world, and a random network (after Watts and Strogatz 1998; reproduced with permission from Science News).

the six degrees of separation (most people are friends with their immediate neighbors but we all have one or two friends a long way away), but it really is not a model found often in the real world. In the real world the architecture of a network is neither random nor small world, but it comes in a variety of distributions, such as truncated power-law distributions (Newmann 2001), Gaussian distributions (Amaral et al. 2000), power-law distributions (Faloutsos et al. 1999), and distributions consisting of two power laws separated by a cutoff value of k (Dorogovtsev and Mendes 2001; Ferrer and Sole 2001). The last two types emerge in certain families of networks that grow in time (Dorogovtsev and Mendes 2001; Barabasi and Albert 1999).

The most interesting and common of such networks are the so-called *scale-free* networks, in which the degree distribution is the power law $p_k \sim k^{-\gamma}$. Consider a map showing an airline’s routes (Fig. 2). This map has a few hubs connected with many other points (supernodes) and many points connected to only a few other points, which is a property associated with power-law distributions. Such a map is highly clustered, yet it allows motion from a point to another faraway point with just a few connections. As such, this network has the *property* of small-world networks, but this property is not achieved by local clustering and a few random connections; it is achieved by having a few elements with a large number of links and many elements with very few links. Thus, even though they share the same property, the architecture of scale-free networks is different than that of small-world networks. Such inhomogeneous networks have been found to pervade biological, social, ecological, and economic systems; the Internet; and other systems (Albert et al. 1999; Jeong et al. 2000; Liljeros et al. 2001; Jeong et al. 2001; Pastor-Satorras and Vespignani 2001; Bouchaud and Mezard 2000; Farkas et al. 2003; Barabasi and Bonabeau 2003; Albert and Barabasi 2002). These networks are referred to as scale free because they show a power-law




FIG. 2. Route map for Continental Airlines (courtesy of Continental Airlines).

distribution of the number of links per node. Lately, it was also shown that, in addition to the power-law-degree distribution, many real scale-free networks consist of self-repeating patterns on all length scales. This result is achieved by the application of a renormalization procedure that coarse grains the system into boxes containing nodes within a given “size” (Song et al. 2005). In other words, scale-free networks also exhibit fractal geometry. These properties are very important because they imply some kind of self-organization within the network. Scale-free networks are not only efficient in transferring information, but, due to the high degree of local clustering, they are also very stable (Barabasi and Bonabeau 2003). Because there are only a few supernodes, chances are that the accidental removal of some nodes will not include the supernodes. In this case the network would not become disconnected. This is not the case with random and, to a lesser degree, small-world networks, where the accidental removal of the same percentage of nodes makes them more prone to failure (Barabasi and Bonabeau 2003; Albert et al. 2000). A scale-free network is vulnerable only when a super-node is “attacked.” Note that scale-free networks have

properties of small-world networks, but small-world networks à la Watts and Strogatz (1998) are not scale free. An example of such a network is given in Fig. 3,




FIG. 3. The network of interactions between the proteins in *Saccharomyces cerevisiae*, otherwise known as baker’s yeast (courtesy of A.-L. Barabasi).

which shows the network of interactions between the proteins in the yeast *Saccharomyces cerevisiae*, otherwise known as baker's yeast (Jeong et al. 2001). By looking at the connectivity of each protein to other proteins, the authors were able to determine that more than 90% of the proteins in the network have less than five links, and that only one in five of these were essential to the survival of the yeast. In other words, removing these proteins did not affect the function of this organism. In contrast, they found that less than 0.7% of the proteins were hubs having many more than 15 connections. For these hubs they found that removal of any hub resulted in the death of the organism. Such findings, which can only be delineated by constructing the network, can be extremely useful because they may lead to ways to protect the organism from microbes by specifically protecting the hubs. In other areas, the presence of scale-free networks has led to strategies to slow the spread of diseases (Lijeros et al. 2001) and secure the Internet (Barabasi and Bonabeau 2003). The networks can be either fixed, where the number of nodes and links remains the same, or evolving, where the network grows as more nodes and links are added (sometimes in the literature growing networks are classified as a new type of network). What-

ever the type of the network, its underlying topology provides clues about the collective dynamics of the network. The basic structural properties of networks are delineated by the clustering coefficient C and the characteristic pathlength (or diameter) L of the network. The clustering coefficient is defined as follows and is illustrated in Fig. 4. Assume that a node i is connected to k_i other nodes. Now, consider the k_i closest nodes of i . This defines the neighborhood of i . Then, count the number of links Δ_i between any two nodes of the neighborhood (excluding node i). The clustering coefficient of node i is then given by $C_i = 2\Delta_i/k_i(k_i - 1)$. Because there can be at most $k_i(k_i - 1)/2$

links between k_i nodes (which will happen if they formed a fully connected subnetwork), the clustering coefficient is normalized on the interval $[0,1]$. The average C_i over all nodes provides C . As such, C provides a measure of local "cliqueness." The diam-

eter of the network is defined by the number of connections in the shortest path between two nodes in the network averaged over all of the pairs of nodes. For a random network having the same average number of connections per node $\langle k \rangle$, it can be shown analytically (Bollabas 1985; Watts and Strogatz 1998; Albert and Barabasi 2002) that $L_{\text{random}} = \ln N / \ln \langle k \rangle$ and $C_{\text{random}} = \langle k \rangle / N$. The small-world property requires that $C \gg C_{\text{random}}$ and $L \geq L_{\text{random}}$. There are other measures and ways to investigate networks. Examples include minimum spanning trees (Mantegna 1999), asset trees and asset graphs (Onnela et al. 2004), tree length and occupation levels (Onnela et al. 2003), and intensity and coherence of networks (Onnela et al. 2005), but here we will stick with the basic principles.

CLIMATE NETWORKS.

How can these ideas be extended to a system like the climate? One way is to assume that interacting dynamical systems can also form a network. Consider, for example, the results shown in Fig. 5. In this figure we start

with a number of limit-cycle (periodic) oscillators with distributed natural frequencies. The state of each oscillator is represented as a dot in the complex plane. The amplitude and phase of each oscillation correspond to the radius and angle in polar coordinates. The color of each oscillator indicates its natural frequency (ranging from violet to red or from high to low frequency). If the oscillators are not coupled, then each oscillator will settle onto its limit cycle and will rotate at its natural frequency. When they are coupled, however, the oscillators appear to self-organize and rotate as a synchronized group with locked amplitudes and phases. When the oscillators




FIG. 4. Illustration of how to estimate the clustering coefficient. (a) A node i is connected to $k_i = 8$ other nodes (solid lines). (b) We consider the $k_i = 8$ closest nodes of i . This defines the neighborhood of i . In this neighborhood there exist $\Delta_i = 4$ connections between nodes (excluding node i) (broken lines). The clustering coefficient of node i is then given by $C_i = 2\Delta_i/k_i(k_i - 1) = 0.143$. The average C_i over all nodes in the network provides the clustering coefficient of the network C .

are more complex than a limit cycle (e.g., chaotic) the situation can be more complicated, but studies have shown that synchronization is possible in this case as well (Strogatz 2001). This synchronization translates into links between the individual oscillators that define the structure of the network of these dynamical systems. Thus, one way to apply networks to the climate system is to assume that climate is represented by a grid of oscillators, with each one representing a dynamical system varying in some complex way. In what we are then interested is the collective behavior of these interacting dynamical systems and the structure of the resulting network. Next, we will present an example that will introduce us to the applications and promise of networks in atmospheric sciences. Some of these ideas have been presented in two recent publications by Tsonis and Roebber (2004) and Tsonis (2004).

We start by considering the global National Centers for Environmental Prediction–National Center for Atmospheric Research (NCEP–NCAR) reanalysis 500-hPa dataset (Kistler et al. 2001). A 500-hPa value indicates the height of the 500-hPa pressure level and provides a good representation of the general circulation (wind flow) of the atmosphere. The data used here are arranged on a grid with a resolution of 5° latitude \times 5° longitude. For each grid point monthly values from 1950 to 2004 are available. This results in 72 points in the east–west direction and 37 points in the north–south direction, for a total of $n = 2,664$ points. These 2,664 points will be assumed to be the nodes of the network. From the monthly values we produced anomaly values (actual value minus the climatological average for each month). Thus, for each grid point we have a time series of 660 anomaly values. In order to define the “connections” between the nodes, the correlation coefficient at lag zero (r) between the time series of all possible pairs of nodes [$n(n - 1)/2 = 3,547,116$ pairs] is estimated. Note that even though r is calculated at zero lag, a connection should not be thought of as being “instantaneous.” The fact that the values are monthly introduces a time scale of at least a month to each connection. Even though most of the annual cycle is removed by producing anomaly val-

ues, some of it is still present because the amplitude of the anomalies is greater in the winter than in the summer. For this reason, in order to avoid spurious high values of r , only the values for December, January, and February in each year were considered. It follows that the estimation of the correlation coefficient between any two time series is based on a sample size of 165. Note that because the values are monthly anomalies there is very little autocorrelation in the time series. A pair is considered as connected if their correlation $|r| \geq 0.5$. This criterion is based on parametric and nonparametric significance tests. According to the Student's t test with $N = 165$, a value of $r = 0.5$ is statistically significant above the 99% level. In addition, randomization experiments where the values of the time series of one node are scrambled and then correlated to the unscrambled values of the time series of the other node indicate that a value of $r = 0.5$ will not arise by chance. The use of the correlation coefficient to define links in networks is not new. Correlation coefficients have been used to successfully derive the topology of gene expression networks (Farkas et al. 2003; de la Fuente et al. 2002; Featherstone and Broadie 2002; Agrawal 2002), and to study financial markets (Mantegna 1999). The choice of $r = 0.5$, while it guarantees statistical significance, is somewhat




FIG. 5. Synchronization of several coupled limit-cycle oscillators. Each oscillator starts from a random initial condition, but soon they all self-organize and rotate as a synchronized group (after Strogatz 2001; reproduced with permission from the author).

arbitrary. The effect of a different correlation threshold is discussed in Tsonis and Roebber (2003). In any case, one may in fact consider all pairs as being connected and study the so-called weighted properties of the network, where each link is assigned a weight proportional to its corresponding correlation coefficient (Onnela et al. 2003, 2004). For the scope of this paper, however, we will keep things simple and consider that a pair is connected if the correlation coefficient is above a threshold.

Once we have decided what constitutes a link, we are ready to look at the architecture of this network and how it relates to dynamics. Figure 6 provides first insight into this question. It shows the area-weighted number of total links (connections) at each geographic location. More accurately, it shows the fraction of the total global area to which a point is connected. This is a more appropriate way to show the architecture of the network because it is a continuous network defined on a sphere, rather than a discrete network on a two-dimensional grid. Thus, if a node i is connected to N other nodes at λ_N latitudes, then its area-weighted connectivity \tilde{C}_i is defined as

$$\tilde{C}_i = \sum_{j=1}^N \cos \lambda_j \Delta A / \sum_{\text{over all } \lambda \text{ and } \phi} \cos \lambda \Delta A, \quad (1)$$

where ΔA is the grid area at the equator and ϕ is the longitude. Once we have the information displayed

in Fig. 6 we can estimate C and L . According to the definition of connectivity [Eq. (1)], in order to find the clustering coefficient of node i , C_i , we consider a circular area on the sphere centered on i that is equal to \tilde{C}_i . Then, C_i is the fraction of this circular area that is connected (for a fully connected area, i.e., all pairs of nodes are connected, $C_i = 1$ for all i). The average C_i over all nodes provides the clustering coefficient of the network C . Note also that according to the definition of \tilde{C} , the average \tilde{C}_i over all nodes gives the clustering coefficient C_{random} . Concerning the estimation of L , rather than finding the number of connections in the shortest path between two points, we estimate the distance of this path on the sphere. For this network we find that $L \cong 9,600$ km and $C = 0.56$. For a random network with the same specifications (number of nodes, and average links per node), it is estimated that $L_{\text{random}} \cong 7,500$ km and $C_{\text{random}} = 0.19$. These values indicate that indeed $L \geq L_{\text{random}}$ and $C > C_{\text{random}}$ (by a factor of 3), which will make this global network close to a small-world network. There is, however, more to this global network than what these values suggest.

Returning to Fig. 6, we observe that it displays two very interesting features. In the Tropics it appears that all nodes possess more or less the same number of connections, which is a characteristic of fully connected networks. In the extratropics it

appears that certain nodes possess more connections than the rest, which is a characteristic of scale-free networks. In the Northern Hemisphere we clearly see the presence of regions where such supernodes exist (in China, North America, and the northeast Pacific Ocean). Similarly, several supernodes are visible in the Southern Hemisphere. These differences between the Tropics and extratropics are clearly delineated in the corresponding degree distributions. Figure 7 shows, on a double-logarithmic plot, the distribution of nodes according to how many links they possess (i.e., p_k against k). Given the definition of a link in




FIG. 6. Total number of links (connections) at each geographic location. The uniformity observed in the Tropics indicates that each node possesses the same number of connections. This is not the case in the extratropics where certain nodes possess more links than the rest.

our case, this figure indicates the fraction of the total area covered as a function of the connectivity \tilde{C} . More specifically, Fig. 7a shows the distribution of nodes in the extratropical region of 30°–65°N and 30°–65°S, and Fig. 7b shows the corresponding distribution of nodes in the Tropics (20°N–20°S). The region from 20° to 30°N and from 20° to 30°S is a transition between the two regimes and was left out for a better delineation of the properties in the Tropics and extratropics. Figure 7a appears to exhibit a scaling regime similar to those observed in scale-free networks. In fact, the slope of this graph is around -2.0 , in agreement with other scale-free networks (Barabasi and Albert 1999). In Fig. 7b no such regime is identifiable. The distribution is basically a narrow peak at about $\tilde{C} = 0.5$, indicating that most points possess the same large number of connections, a characteristic of regular, almost fully connected networks. Deviations from the power law (manifested as a peak at about $\tilde{C} = 0.4$ in Fig. 7a) and uniformity (the four points below $\tilde{C} = 0.1$ in Fig. 7b) are due to the fact that the two subnetworks are interwoven; a node in one subnetwork may be connected to a node or nodes in the other subnetwork. Note that very similar results are obtained if, instead of the 30°–65°N and 30°–65°S belts, the whole extratropical area (35°N–90° and 35°S–90°) is considered. It thus appears that the overall network is a “fusion” of a fully connected tropical network and a scale-free extratropical network. As is the case with all scale-free networks, the extratropical subnetwork is also a small-world network. Indeed, we find that for points in the extratropics, the clustering coefficient is much greater than that of a corresponding random network (by a factor of 9). The collective behavior of the individual dynamical systems in the complete network is not described by a single type, but it self-organizes into two coupled subnetworks—one regular, almost fully connected network, operating in the Tropics, and one scale-free/small-world network, operating in the higher latitudes. The extratropics are considered as one subsystem, even though they are physically separated. Whether or not we consider them as one or two subsystems does not modify the physical interpretation, which is that the equatorial network acts as an agent that connects the two hemispheres, thus allowing information to flow between them. This interpretation is consistent with the various suggested mechanisms for interhemispheric teleconnections (Tomas and Webster 1994; Love 1985; Compo et al. 1999; Meehl et al. 1996; Carrera 2001), with the notion of subsystems in climate proposed in




FIG. 7. Degree distribution of (a) extratropical (30°–65°N, 30°–65°S) and (b) tropical (20°N–20°S) grid points.

the late 1980s (Tsonis and Elsner 1989; Lorenz 1991), and with recent studies on synchronized chaos in the climate system (Duane et al. 1999).

An interesting observation in Fig. 6 is that supernodes may be associated with major teleconnection patterns. For example, the supernodes in North America and the northeast Pacific Ocean coincide with the well-known Pacific–North America (PNA) pattern (Wallace and Gutzler 1981). In the Southern Hemisphere we also see supernodes over the southern tip of South America, Antarctica, and the south Indian Ocean that are consistent with some of the features of the Pacific–South America (PSA) pattern (Mo and Higgins 1998). Interestingly, no such supernodes are evident where the other major pattern, the North Atlantic Oscillation (NAO; Thompson and Wallace 1998; Pozo-Vazquez et al. 2001; Huang et al. 1998), is found. This does not indicate that NAO is not

a significant feature of the climate system. Because NAO is not strongly connected to the Tropics, the high connectivity of the Tropics with other regions masks NAO out. In fact, if we consider a network with only nodes north of 30°N latitude, we find (Fig. 8) that a dipole consistent with NAO is not only present, but is also a prominent feature of the network. It should be noted here that in their pioneering paper Wallace and Gutzler (1981) defined teleconnectivity at each grid point as the strongest negative correlation between a grid point and all other points. This brings out teleconnection patterns associated with waves, such as the trough–ridge–trough PNA pattern. However, because of the requirement of the strongest negative correlation (which occurs between a negative anomaly center and a positive anomaly center), this approach can only delineate long-range connections. As such, information about clustering and connectivity at other spatial scales is lost. In the network approach all of the links at a point are considered, and as such much more information (clustering coefficients, diameter, scaling properties, etc.) can be obtained. The similarities between Wallace and Gutzler’s results and the network results arise from the fact that grid points with many long-range links will most likely stand out.

The physical interpretation of the results is that the climate system (as represented by the 500-hPa field) exhibits properties of stable networks and networks where information is transferred efficiently. In the case of the climate system, “information” should be regarded as “fluctuations” from any source. These fluctuations will tend to destabilize the source region. For example, dynamical connections between the ocean and the atmosphere during El Niño may make the

climate over the tropical Pacific less stable. However, the small-world as well as the scale-free properties of the extratropical network and the fully connected tropical network allow the system to respond quickly and coherently to fluctuations introduced into the system. This information transfer diffuses local fluctuations, thereby reducing the possibility of prolonged local extremes and providing greater stability for the global climate system. An important consequence of this property is that local events may have global implications. The fact that the climate system may be inherently stable may not come as a surprise to some, but it is interesting that we find that a stable climate may require teleconnection patterns.

Unlike networks where a node is solidly defined (think of social networks where a node is a person), here a node is a point on a grid, which is defined rather arbitrarily and/or can be represented at various resolutions. Strictly speaking, our network has infinite nodes. Due, however, to spatial correlations, the “effective” number of nodes is much less. In fact, we find that reproducing Fig. 6, but with the full grid of 2.5° × 2.5° resolution, results in virtually the same network architecture. Other studies have demonstrated this as well. For example, in the *www* network an increase in the number of nodes by a factor of 2,500 results in an increase of *L* by a factor of only 1.6, and the architecture (vis-à-vis degree distribution) remains identical (Albert and Barabasi 2002). Other meteorological fields may also exhibit small-world or scale-free properties. As an additional example we used an upper-tropospheric streamfunction. Figure 9 is similar to Fig. 8, but for the streamfunction. Again, here we observe supernodes. This network is also a scale-free and a small-world network.




FIG. 8. Same as Fig. 6, but we start with a network with nodes north of 30°N latitude only.

DISCUSSION. From this initial application of networks to climate it appears that atmospheric fields can be thought of as a network of interacting points whose collective behavior may exhibit properties of small-world networks. This ensures the efficient transfer of information. In addition, the scale-free architectures guarantee stability. Furthermore, supernodes in the network identify teleconnection patterns. As was demonstrated in Tsonis (2004), these teleconnections are not static phenomena, but their spatiotemporal variability is affected by large (global) changes. The 55-yr period used to produce Fig. 6 can be divided into two distinct periods, each with a length of 27 yr (1951–77 and 1978–2004). During the first period the global temperature shows no significant overall trend. During the second period, however, a very strong positive trend is present. Because there is a distinct




FIG. 9. Same as Fig. 8, but for the streamfunction.

change in the global property of the system, how does this affect the dynamics of the global network? To answer this question C and L for the two periods were estimated. It was found that C is about 5% smaller and L is about 4% smaller in the second period. This result will indicate that during the warming of the planet the network has acquired more long-range and less small-range connections. This is clearly shown in Fig. 10, which shows the distribution of connections according to their distance (as calculated on the sphere). The solid line represents the distribution in the first period and the dashed line represents the distribution in the second period. This figure shows that the frequency of the long-range connections ($>6,000$ km) has increased, whereas the frequency of the shorter-range connections ($<6,000$ km) has decreased. The differences between the two distributions may not appear to be impressive, but with hundreds of thousands of connections involved, these differences are, according to the Kolmogorov–Smirnov test, statistically significant at the 99% confidence level (see also Tsonis 2004). Even though this is only one example, this result suggests that monitoring the properties of such networks may provide an additional tool to identify or verify climate changes.

Furthermore, mapping atmospheric fields into networks appears to bring out properties of the general circulation. Thus, it may provide an alternative approach to study atmospheric phenomena and dynamics. Moreover, just because in the case of 500-hPa teleconnections the network approach brings out that which has been found by the linear approaches (such as EOF analysis), it does not mean that it will always produce what the linear approaches produce. For example, scale-free phenomena are associated with

nonlinear dynamics. As such, linear approaches, such as EOF analysis, *cannot* bring out this property. The fact that our network approach recovers the scale-free characteristic is a strong indication that it will *not* always produce the same result as a linear approach, and that in fact it may produce novel insights. The following presents another example where linear approaches would not yield certain properties. The current standard for understanding nonlocal interactions in the atmosphere is linear inverse modeling (e.g., Winkler et al. 2001). In plain terms, linear inverse modeling is similar to a least squares fit linking the values of certain dynamical quantities at one time (e.g., a subset of empirical orthogonal functions) with their values at some future time. In practice, this involves making the assumption that the dynamics are sufficiently approximated by a linear, stable, stochastic dynamical system, and then the propagator for that system is calculated from data at some fixed time lag. It is not clear that such linear approaches are necessarily optimal, however, and specifically, whether they distort the network structure of the atmosphere. For an example of such distortion, consider a one-dimensional discrete dynamical system, similar to that introduced by Lorenz and Emanuel (1998), that is chaotic and mimics zonal wave propagation in the atmosphere. The relevant time scale in this system is the error-doubling time, which we take as one model day. This model has 40 nodes, and is modified to possess a long-range spatial correlation by allowing nodes 20 and 40 to force each other. Given




FIG. 10. The relative frequency distribution of the connections according to their distance for the periods of 1951–77 (solid line) and 1978–2004 (dashed line). This result indicates that during periods of warming, the network acquires more long-range connections and less small-range connections.

this setup, we calculated the correlation for each node with every other node at a lag of five model days and defined links for $|r| \geq 0.5$. Our rationale in doing this was to see how effectively information is transferred into the future. The solid line in Fig. 11 shows the number of links of each node. We observe that most of the nodes possess just about four to five links, but a few nodes stand above this level with more links. Interestingly, as expected from a nonlinear system, it is not necessarily nodes 20 and 40 that possess most of the links. However, a linear inverse model (broken lines) constructed to provide a 5-day forecast does not possess a similar connectivity pattern. Apparently, the linear inverse approach does not preserve the properties of the actual nonlinear model.

True enough, the correlation coefficient is a linear measure. One may question why we should use a linear measure to study nonlinear dynamics. This is a legitimate question. A nonlinear measure that could be used instead is the mutual information, but the problem is that its accurate estimation requires much more data than we have available. As such, there is very little choice but to use the correlation coefficient as an indicator of a link. However, the correlation coefficient is used only as a construction tool. In a similar way it has been used successfully in the past to delineate nonlinear dynamics. For example, in reconstructing attractors from time series, the state space is reconstructed using coordinates that are shifts of the original time series. The optimum shift is often assumed to be the lag at which the autocorrelation function first becomes zero. This approach reproduces the properties of the attractor quite reasonably (Tsonis 1992; Tsonis and Elsner 1989).




FIG. 11. An example demonstrating that linear approaches may not yield the true result of a nonlinear process.

OUTLOOK. Complex networks describe many natural and social dynamical systems, and their study has revealed interesting mechanisms underlying their function. The novelty of networks is that they bring out topological/geometrical aspects that are related to the physics of the dynamical system in question, thus providing a new and innovative way to treat and investigate nonlinear systems and data. While several advances have been made, this area is still young and the future is wide open. This introductory paper presented some fundamental aspects of networks and some preliminary results of the application of networks to climatic data, which indicate that networks delineate some key features of the climate system. This suggests that networks have the potential to become a new and useful tool in climate research.

REFERENCES

- Agrawal, H., 2002: Extreme self-organization in networks constructed from gene expression data. *Phys. Rev. Lett.*, **89**, 268–702.
- Albert, R., and A.-L. Barabasi, 2002: Statistical mechanics of complex networks. *Rev. Mod. Phys.*, **74**, 47–101.
- , H. Jeong, and A.-L. Barabasi, 1999: Diameter of the World Wide Web. *Nature*, **401**, 130–131.
- , —, and —, 2000: Error and attack tolerance of complex networks. *Nature*, **406**, 378–382.
- Amaral, L. A. N., A. Scala, M. Barthelemy, and H. E. Stanley, 2000: Classes of behavior of small world networks. *Proc. Natl. Acad. Sci. USA*, **97**, 11 149–11 152.
- Barabasi, A.-L., and R. Albert, 1999: Emergence of scaling in random networks. *Science*, **286**, 509–512.
- , and E. Bonabeau, 2003: Scale-free networks. *Sci. Amer.*, **288**, 60–69.
- Bollabas, B., 2001: *Random Graphs*. 2d ed. Cambridge University Press, 498 pp.
- Bouchaud, J.-P., and M. Mezard, 2000: Wealth condensation in a simple model of economy. *Physica A*, **282**, 536–540.
- Carrera, M. L., 2001: Significant events of inter-hemispheric atmospheric mass exchange. Ph.D. thesis, McGill University, 172 pp.
- Compo, G. P., G. N. Kiladis, and P. J. Webster, 1999: The horizontal and vertical structure of east Asian winter monsoon pressure surges. *Quart. J. Roy. Meteor. Soc.*, **125**, 29–54.
- de la Fuente, A., P. Brazhnik, and P. Mendes, 2002: Linking the Genes: Inferring quantitative gene networks from microarray data. *Trends Genet.*, **18**, 395–398.
- Dorogovtsev, S. N., and J. F. F. Mendes, 2001: Language as an evolving word web. *Proc. Roy. Soc. London B*, **268**, 2603–2602.

- Duane, G. S., P. J. Webster, and J. B. Weiss, 1999: Co-occurrence of Northern and Southern Hemisphere blocks as partially synchronized chaos. *J. Atmos. Sci.*, **56**, 4183–4205.
- Erdos, P., and A. Renyi, 1960: On the evolution of random graphs. *Publ. Math. Inst. Hung. Acad. Sci.*, **5**, 17–61.
- Faloutsos, M., P. Faloutsos, and C. Faloutsos, 1999: On power-law relationships of the internet topology. *Comput. Comm. Rev.*, **29**, 251–260.
- Farkas, I. J., H. Jeong, T. Vicsek, A.-L. Barabási, and Z. N. Oltvai, 2003: The topology of the transcription regulatory network in the yeast *Saccharomyces cerevisiae*. *Physics*, **318A**, 601–612.
- Featherstone, D. E., and K. Broadie, 2002: Wrestling with pleiotropy: Genomic and topological analysis of the yeast gene expression network. *Bioessays*, **24**, 267–274.
- Ferrer, R., and R. V. Sole, 2001: The small-world of human language. *Proc. Roy. Soc. London B*, **268**, 2261–2266.
- Huang, J. P., K. Higuchi, and A. Shabbar, 1998: The relationship between the North Atlantic Oscillation and El Niño Southern Oscillation. *Geophys. Res. Lett.*, **25**, 2707–2710.
- Jeong, H., B. Tombor, R. Albert, A. N. Oltvai, and A.-L. Barabasi, 2000: The large scale organization of metabolic networks. *Nature*, **407**, 651–654.
- , S. Mason, A.-L. Barabasi, and Z. N. Oltvai, 2001: Lethability and centrality in protein networks. *Nature*, **411**, 41–42.
- Kistler, R., and Coauthors, 2001: The NCEP–NCAR 50-year reanalysis: Monthly means, CD-ROM, and documentation. *Bull. Amer. Meteor. Soc.*, **82**, 247–267.
- Liljeros, F., C. Edling, L. N. Amaral, H. E. Stanley, and Y. Aberg, 2001: The web of human sexual contacts. *Nature*, **411**, 907–908.
- Lorenz, E. N., 1991: Dimension of weather and climate attractors. *Nature*, **353**, 241–244.
- , and K. A. Emanuel, 1998: Optimal sites for supplementary weather observations: Simulation with a small model. *J. Atmos. Sci.*, **55**, 399–414.
- Love, G., 1985: Cross-equatorial influence of winter hemisphere subtropical cold surges. *Mon. Wea. Rev.*, **113**, 1487–1498.
- Mantegna, R. N., 1999: Hierarchical structure in financial markets. *Eur. Phys. J.*, **11B**, 193–197.
- Meehl, G. A., G. N. Kiladis, K. M. Weickmann, M. Wheeler, D. S. Gutzler, and G. P. Compo, 1996: Modulation of equatorial subseasonal convective episodes by tropical-extratropical interaction in the Indian and Pacific Ocean regions. *J. Geophys. Res.*, **101**, 15 033–15 049.
- Milgram, S., 1967: The small-world problem. *Psych. Today*, **1**, 60–67.
- Mo, K. C., and R. W. Higgins, 1998: The Pacific-South America modes and tropical convection during the Southern Hemisphere winter. *Mon. Wea. Rev.*, **126**, 1581–1596.
- Newmann, M. E. J., 2001: The structure of scientific collaboration networks. *Proc. Natl. Acad. Sci. USA*, **98**, 404–409.
- Onnela, J.-P., A. Chakraborti, K. Kaski, J. Kertesz, and A. Kanto, 2003: Dynamics of market correlations: Taxonomy and portfolio analysis. *Phys. Rev. E*, **68**, doi:10.1103/PhysRevE.68.056110.
- , K. Kaski, and J. Kertesz, 2004: Clustering and information in correlation based financial networks. *Eur. Phys. J.*, **38B**, 353–362.
- , J. Saramaki, J. Kertesz, and K. Kaski, 2005: Intensity and coherence of motifs in weighted complex networks. *Phys. Rev. E.*, **71**, doi:10.1103/PhysRevE.71.065103.
- Pastor-Satorras, R., and A. Vespignani, 2001: Epidemic spreading in scale-free networks. *Phys. Rev. Lett.*, **86**, 3200–3203.
- Pozo-Vazquez, D., M. J. Esteban-Parra, F. S. Rodrigo, and Y. Castro-Diez, 2001: The association between ENSO and winter atmospheric circulation and temperature in the North Atlantic region. *J. Climate*, **14**, 3408–3420.
- Song, C., S. Havlin, and H. A. Makse, 2005: Self-similarity of complex networks. *Nature*, **433**, 392–395.
- Strogatz, S. H., 2001: Exploring complex networks. *Nature*, **410**, 268–276.
- Thompson, D. W. J., and J. M. Wallace, 1998: The Arctic Oscillation signature in the wintertime geopotential height and temperature fields. *Geophys. Res. Lett.*, **25**, 1297–1300.
- Tomas, R. A., and P. J. Webster, 1994: Horizontal and vertical structure of cross-equatorial wave propagation. *J. Atmos. Sci.*, **51**, 1417–1430.
- Tsonis, A. A., 1992: *Chaos: From Theory to Applications*. Plenum Press, 274 pp.
- , 2004: Does global warming inject randomness into the climate system? *Eos, Trans. Amer. Geophys. Union*, **85**, 361–364.
- , and J. B. Elsner, 1989: Chaos, strange attractors, and weather. *Bull. Amer. Meteor. Soc.*, **70**, 16–23.
- , and P. J. Roebber, 2004: The architecture of the climate network. *Physica*, **333A**, 497–504.
- Wallace, J. M., and D. S. Gutzler, 1981: Teleconnections in the geopotential height field during the Northern Hemisphere winter. *Mon. Wea. Rev.*, **109**, 784–812.
- Watts, D. J., and S. H. Strogatz, 1998: Collective dynamics of “small-world” networks. *Nature*, **393**, 440–442.
- Winkler, C. R., M. Newman, and P. D. Sardeshmukh, 2001: A linear model of wintertime low-frequency variability. Part I: Formulation of forecast skill. *J. Climate*, **14**, 4474–4494.