

APOE4 homozygote females are resistant to the beneficial effects of 17 β -estradiol on memory and CA1 dendritic spine density in the EFAD mouse model of Alzheimer's disease

Lisa R. Taxier^a, Sarah M. Philippi^{a,c}, Aaron W. Fleischer^a, Jason M. York^b, Mary Jo LaDu^b, Karyn M. Frick^{a,*}

^a Department of Psychology, University of Wisconsin-Milwaukee, Milwaukee WI, USA

^b Department of Anatomy and Cell Biology, University of Illinois at Chicago, Chicago, IL, USA

^c Current affiliation: Department of Neuroscience and Neuroscience Program, Icahn School of Medicine at Mount Sinai, New York, NY, USA



ARTICLE INFO

Article history:

Received 30 December 2021

Revised 1 June 2022

Accepted 17 June 2022

Available online 23 June 2022

Keywords:

Alzheimer's disease

Apolipoprotein E

APOE4

17 β -estradiol

Spatial memory

Object recognition

Hippocampus

Medial prefrontal cortex

ABSTRACT

Female *APOE4* carriers are at greatest risk of Alzheimer's disease (AD). The potent estrogen 17 β -estradiol (E_2) may mediate AD risk, as the onset of memory decline coincides with the menopausal transition. Whether *APOE* genotype mediates E_2 's effects on memory and neuronal morphology is poorly understood. We used the *APOE^{+/+}/5xFAD^{+/−}* (EFAD) mouse model to examine how *APOE3* homozygote (E3FAD), *APOE3/4* heterozygote (E3/4FAD), and *APOE4* homozygote (E4FAD) genotypes modulate effects of E_2 on object and spatial memory consolidation, dendritic spine density, and dorsal hippocampal estrogen receptor expression in 6-month-old ovariectomized EFAD mice. Dorsal hippocampal E_2 infusion enhanced memory consolidation and increased CA1 apical spine density in E3FAD and E3/4FAD, but not E4FAD, mice. CA1 basal mushroom spines were also increased by E_2 in E3FADs. E4FAD mice exhibited reduced CA1 and mPFC basal spine density, and increased dorsal hippocampal ER α protein, independent of E_2 . Overall, E_2 benefitted hippocampal memory and structural plasticity in females bearing one or no *APOE4* allele, whereas two *APOE4* alleles impeded the memory-enhancing and spinogenic effects of E_2 .

© 2022 Elsevier Inc. All rights reserved.

1. Introduction

Alzheimer's disease (AD), the most common form of dementia, poses a monumental global health crisis. Despite efforts by researchers to identify pharmacological treatments and drug targets, there are few disease-modifying therapies that mitigate the devastating impact of AD. Other than aging, risk factors for AD include *APOE4* genotype and female sex. *APOE4* genotype renders individuals more susceptible to AD than other *APOE* genotypes, increasing risk by as much as 2-fold in the presence of a single *APOE4* allele (Bertram, 2009; Roses, 1996). Women, especially *APOE4* car-

riers, are at significantly greater risk for developing AD compared to men (Altmann et al., 2014; Bretsky et al., 1999), making the examination of factors contributing to this increased risk in women critical for AD research.

One potential mediator of heightened risk for AD in women is the loss of circulating estrogens coincident with the onset of the menopausal transition. Multiple studies implicate the precipitous loss of 17 β -estradiol (E_2), the most prevalent circulating estrogen, at menopause as a contributing factor for age and disease-associated cognitive decline (Jacobs et al., 2016; Paganini-Hill and Henderson, 1996). In fact, female AD patients have lower levels of endogenous E_2 than age and sex-matched controls (Manly et al., 2000; Tsolaki et al., 2005). However, clinical trials of estrogen therapy paint a conflicting portrait of its effects on AD risk; early studies offered support for the idea that E_2 may reduce risk of AD (Baldereschi et al., 1998; Joffe et al., 2006; Kawas et al., 1997; Tang et al., 1996), whereas other studies and more recent clinical trials do not (Gleason et al., 2015; Rapp et al., 2003; Shumaker et al., 2003). These discrepant reports may be related to key factors influencing estrogen therapy efficacy including age at initiation, specific estrogen treatment regimen (e.g., estradiol vs.

Abbreviations: 5XFAD, 5 familial AD mutations; AD, Alzheimer's Disease; ANOVA, Analysis of Variance; DH, Dorsal Hippocampus; E_2 , 17 β -estradiol; EFAD, *APOE^{+/+}/5xFAD^{+/−}*; E3FAD, *APOE^{3/3}/5xFAD^{+/−}*; E3/4FAD, *APOE^{3/4}/5xFAD^{+/−}*; E4FAD, *APOE^{4/4}/5xFAD^{+/−}*; ER α , estrogen receptor alpha; ER β , estrogen receptor beta; GPER, G-protein-coupled estrogen receptor; mPFC, medial prefrontal cortex; OP, Object Placement; OR, Object Recognition; Tg, Transgenic; S, Second; Min, Minute; H, Hour.

* Corresponding author at: Department of Psychology, University of Wisconsin-Milwaukee, 2441 E. Hartford Ave., Milwaukee, WI, 53211.

E-mail address: frick@uwm.edu (K.M. Frick).

conjugated equine estrogens), dose, length of treatment, and physical health (Henderson, 2006; Paganini-Hill and Henderson, 1996). Another critical element of cognitive responsiveness to estrogen therapy is *APOE* status. Importantly, cognition in women *APOE4* carriers appears to benefit less from estrogen therapy than in their *APOE4*-negative counterparts (Yaffe et al., 2000). This clinical work is consistent with evidence from basic research; *APOE2* and *APOE3*, but not *APOE4*, act synergistically with E_2 to promote neurite outgrowth in mixed cell cultures, suggesting that *APOE4* expression renders cells uniquely nonresponsive to E_2 (Nathan et al., 2004). Combined, the above results suggest that E_2 may be beneficial for carriers of *APOE3* alleles, and unhelpful or even deleterious for *APOE4* carriers.

In adult female mice, treatment with exogenous E_2 facilitates memory via rapid activation of cell signaling within the dorsal hippocampus (DH), which is accompanied by an increase in dendritic spine density in the DH and in the medial prefrontal cortex (Boulware et al., 2013; Fernandez et al., 2008; Fortress et al., 2013; Tuscher et al., 2016a). The spinogenic effects of E_2 have been linked to improvements in learning and memory (Inagaki et al., 2012; Phan et al., 2015; Smith et al., 2009), suggesting that spinogenesis may be used as a morphological biomarker of the pro-cognitive effects of E_2 . Importantly, spine density loss in the hippocampus and cortex is an established correlate of cognitive decline in AD patients, and in some cases is associated more strongly with aberrant memory function than other pathological hallmarks of AD (DeKosky and Scheff, 1990; Selkoe, 2002). Unsurprisingly, *APOE4* has deleterious effects on spine density relative to *APOE3* in both transgenic mice and in humans, which may drive accelerated disease progression (Dumanis et al., 2009; Ji et al., 2003). Given that synaptic integrity and dendritic spine density are closely linked to learning and memory, spinogenic compounds, such as E_2 , may be of benefit to AD patients. However, whether the spinogenic or mnemonic benefits of E_2 are modulated by *APOE* genotype is unknown.

The present study was designed to test whether *APOE* genotype interacts with E_2 to modulate multiple forms of memory, dendritic spine density in the DH and medial prefrontal cortex (mPFC), and DH cell signaling and estrogen receptor expression in the well-characterized EFAD mouse model of Alzheimer's disease (Liu et al., 2015; Tai et al., 2017; Youmans et al., 2012). Female EFAD-transgenic (Tg) mice (*APOE*^{+/+}/5xFAD^{+/-}) express 5 familial AD (FAD) mutations (5xFAD), and human *APOE3* (*APOE*^{3/3}/5xFAD^{+/-}; E3FAD), *APOE3/4* (*APOE*^{3/4}/5xFAD^{+/-}; E3/4FAD), or *APOE4* (*APOE*^{4/4}/5xFAD^{+/-}; E4FAD). At 6 months of age, the hippocampus and frontal cortex of E4FAD mice express striking AD-like pathology, with significantly higher accumulation of total $\text{A}\beta42$, soluble $\text{A}\beta42$, soluble oligomeric $\text{A}\beta42$, amyloid plaques, and total *APOE* relative to E3FAD mice (Youmans et al., 2012). Thus, this model allows examination of the influence of *APOE* genotype on the cognitive and neural response to E_2 against a background of AD-like pathology, which makes the EFAD model ideal for this study. First, ovariectomized female E3FAD, E3/4FAD, and E4FAD mice received bilateral DH infusion of E_2 immediately after training in object recognition (OR) and object placement (OP) tasks. Given the vulnerability of the DH and mPFC to excitatory synapse loss in AD, we next quantified dendritic spine density in the CA1 region of the DH and the mPFC. Because previous studies in wild-type mice showed that E_2 rapidly increases activation of estrogen receptors and numerous cell-signaling kinases in the DH (e.g., Boulware et al., 2013; Kim et al., 2016, 2019), we also used western blotting to assess DH levels of kinase activity, estrogen receptor alpha (ER α), estrogen receptor beta (ER β), and G-protein-coupled estrogen receptor (GPER). We hypothesized that E3FADs would be the

most responsive to E_2 , such that DH E_2 infusion would promote memory consolidation in both tasks and would increase dendritic spine density in both the DH and mPFC. We also hypothesized that E4FADs would be the least responsive to E_2 , and that E_2 's efficacy in E3/4FADs would be intermediate between E3FADs and E4FADs. Based on our previous study of sex differences in E3FAD and E4FAD mice (Taxier et al., 2022a), we expected estrogen receptor alpha (ER α) levels to be highest in the DH of E4FAD mice but predicted effects on other estrogen receptors were unclear based on a lack of previous data on expression of these receptors in EFAD mice. We also anticipated that mnemonic benefits of E_2 in EFADs, if present, could be linked to rapid activation of cell signaling. We found that DH infusion of E_2 facilitated memory consolidation and an increase in CA1 apical dendritic spine density in ovariectomized E3FAD and E3/4FAD, but not E4FAD, mice. Thus, in mice bearing significant AD-like pathology, E_2 benefitted memory and structural plasticity in mice with 0 or 1 copy of *APOE4*, whereas 2 copies of *APOE4* blocked the memory-enhancing and spinogenic effects of E_2 . Understanding how *APOE4* impedes the actions of E_2 may provide new insights into the variable effects of E_2 on AD risk in women.

2. Methods

2.1. Subjects

APOE-TR^{+/+}/5xFAD^{+/-} (EFAD) mice express 5 familial AD mutations (APP K670N/M671L, I716V, V717I, PS1 M146L, and L286V) under control of the neuron-specific mouse Thy-1 promoter, and express human *APOE3* or *APOE4* (Tai et al., 2017; Youmans et al., 2012). Female mice for the present experiment were homozygous for human *APOE3* (E3FAD) or *APOE4* (E4FAD), or had 1 copy each of *APOE3* and *APOE4* (E3/4FAD). EFAD mice were bred, weaned, and genotyped at the University of Illinois at Chicago (UIC; Animal use protocol 17-066) and shipped to the University of Wisconsin-Milwaukee (UWM; Animal use protocol 19-20-03) at 2 months of age, where they were aged to 6 months before the start of behavioral testing. Prior to surgery, mice were housed in groups of up to 5 per cage and were singly housed following cannulation and ovariectomy. Mice were behaviorally tested in 4 separate cohorts, whereas brain analyses for all mice were conducted at the same time. The first 2 behavioral cohorts consisted of only E3FAD and E4FAD genotypes because E3/4FAD mice were not available at that time. When sufficient numbers of E3/4FAD mice could be generated for this project, they were added to the latter 2 cohorts, which included mice from all 3 genotypes. Thus, Ns for the E3/4FAD groups are smaller than those of the E3FAD and E4FAD groups. A subset of brains was dissected and frozen for western blotting and the rest were Golgi impregnated for spine density analyses. Mice were maintained on a 12hour light/dark cycle with *ad libitum* access to food and water for the duration of the study. Protocols and procedures followed the National Institutes of Health Guide for the Care and Use of Laboratory Animals and were approved by the University of Wisconsin-Milwaukee Institutional Animal Care and Use Committee.

2.2. Surgeries

At the start of surgery, mice received a 5 mg/kg dose of subcutaneous Rimadyl for pain management. They were then anesthetized with isoflurane in 100% oxygen (5% for induction, 2% for maintenance) and placed in a stereotaxic apparatus (Kopf instruments) for ovariectomy and cannula implantation, which occurred within the same surgical session (Lewis et al., 2008; Taxier et al., 2019). Stainless steel bilateral guide cannulae (C232GC, 22 gauge;

Plastics One) were implanted into the DH (-1.7 mm AP, \pm 1.5 mm ML, and -2.3 mm DV) and affixed to the skull using dental cement (Darby Dental). Dummy cannulae (C232DC; Plastics One) were used to prevent clogging of the guide cannulae. Mice received MediGel carprofen (ClearH2O) for postsurgical analgesia and were given one week to recover prior to the start of behavioral testing.

2.3. Drugs and infusions

Hydroxypropyl- β -cyclodextrin (HBC)-encapsulated E₂ (Sigma-Aldrich) was dissolved in 0.9% sterile saline to a concentration of 10 μ g/ μ l and was infused at a rate of 0.5 μ l/min for 1 min/hemisphere (Fortress et al., 2013; Kim et al., 2019; Taxier et al., 2019). Vehicle-treated mice received HBC (Sigma-Aldrich) dissolved in saline to the same concentration of cyclodextrin as in the E₂ solution. For behavioral experiments, mice received a bilateral DH infusion of vehicle or E₂ immediately post-training. Two weeks separated bouts of behavioral training to ensure that any acute effects of post-training infusions had dissipated prior to the next infusion. Two weeks following the conclusion of behavioral testing, mice were reinfused with the same treatment they received previously, and tissue was collected 5 minutes later for western blotting or 2 hours later for Golgi impregnation and spine counting.

2.4. Behavioral tasks

Object recognition (OR) and object placement (OP) were conducted as described previously (Boulware et al., 2013; Fernandez et al., 2008; Fortress et al., 2013; Kim et al., 2019; Taxier et al., 2019; Tuscher et al., 2018). Mice were handled for 30 s/day for 3 days prior to exposure to the behavioral arena. On the second day of handling, a Lego Duplo block (6.3 \times 3.1 \times 2.3 cm) was placed into each home cage to acclimate mice to objects for the remainder of handling and habituation. After handling was completed, mice were allowed to habituate to the empty behavioral arena (60 cm \times 60 cm \times 47 cm) for 5 min/d for 2 days. Light levels in the area were measured daily to ensure even illumination of 75 \pm 5 lux in each corner of the arena. The following day, mice were acclimated to the empty arena for 2 minutes, removed to a holding cage, and immediately returned to the arena and exposed to 2 identical objects placed 5 cm from the upper left and right corners of the arena. Mice were then allowed to freely explore the objects until they accumulated 30 seconds of object exploration (or until 20 minutes had elapsed), defined as any time a mouse's nose was within 2 cm of the object. The 30 second criterion was chosen to ensure that mice spent an equivalent amount of time exploring the training objects prior to drug infusion, as different amounts of time spent with these objects could alter the strength of the memory in such a way as to occlude effects of drug manipulation. Mice that did not reach the 30 second criterion within the allotted 20 minutes were excluded. Immediately following training, mice received bilateral DH infusion of HBC vehicle or E₂ and were then returned to their home cage.

OR testing was conducted 24 hours after drug infusion, at which point mice were returned to the arena in which one familiar object from the training phase was replaced by a novel object. Mice were again allowed to accumulate 30 seconds of object exploration. OP testing was conducted 4 hours after drug infusion; here, mice were allowed to explore one familiar training object in its original location, and the second training object in a new location (bottom left or right corner of the arena). Mice that remember the identity and location of the training objects, respectively, spend more time than chance (15 second) exploring the novel or moved objects during testing (Kim et al., 2019; Taxier et al., 2019; Tuscher et al., 2016b). Time spent with the

objects and time to accumulate 30 seconds of exploration were recorded using ANYmaze automated tracking software and scored by a researcher blinded to treatment and genotype (San Diego Instruments). The order of OR and OP testing was counterbalanced across groups, and approximately 2 weeks separated OR training and testing from OP training and testing to allow for acute effects of post-training E₂ infusion to dissipate prior to the next training session.

Although vehicle-treated ovariectomized wild-type mice typically remember the identity and location of the training objects after a 24 hours (OR) or 4 hours (OP) delay, respectively (Kim et al., 2019; Taxier et al., 2019; Tuscher et al., 2016b), we have previously shown that only male E3FADs, and not female E3FADs or E4FADs of either sex, remember the training object identity or location when tested at these same delays (Taxier et al., 2022a). Thus, given that female E3FADs and E4FADs exhibit impaired memory at the 24 hours (OR) and 4 hours (OP) delays, mice in the present experiment were tested using these delays to assess whether DH E₂ infusion could facilitate memory consolidation.

2.5. Western blotting

Two weeks after the conclusion of behavioral testing, a subset of mice (10–12/group) were infused into the DH with vehicle or E₂. Mice were cervically dislocated and decapitated 5 minutes later, based on previous studies in wild-type mice showing that E₂ or GPER activation increases phosphorylation of numerous cell-signaling kinases, including ERK and JNK, in the DH at this time point (e.g., Boulware et al., 2013; Kim et al., 2016, 2019). Brains from each group were extracted, and the DH was bilaterally dissected immediately on ice and frozen at -80°C. Western blotting was conducted as described previously (Boulware et al., 2013; Fernandez et al., 2008; Kim et al., 2019; Koss et al., 2018; Taxier et al., 2019). Tissue samples were resuspended 1:25 weight/volume in lysis buffer containing PMSF and a protease inhibitor cocktail (Sigma-Aldrich) and homogenized using a probe sonicator (Branson Sonifier 250). Homogenates were electrophoresed on 10% TGX (Tris-Glycine eXtended) stain-free precast gels (Bio-Rad) and transferred to polyvinylidene fluoride (PVDF) membranes (Bio-Rad) using the TransBlot Turbo system (Bio-Rad). Total protein transfer was then verified using a ChemiDoc MP gel imager (Bio-Rad).

Membranes were blocked in 5% milk and incubated with following primary antibodies overnight at 4°C: phosphorylated 42/44 ERK (#9101, 1:2000, Cell Signaling Technology), total ERK (#9102, 1:20000, Cell Signaling Technology), phosphorylated 46/54 JNK (#4668, 1:1000, Cell Signaling Technology), total JNK (#9252, 1:1000, Cell Signaling Technology), ER α (H-184, 1:1000, Santa Cruz Biotechnology), ER β (#PA1-310B, 0.25 μ g/mL, Thermo Fisher), and GPER (ab39742, 1:250, Abcam). The following day, all blots were incubated at room temperature with a rabbit (#7074, 1:5000, Cell Signaling Technology) or mouse (#7076, 1:5000, Cell Signaling Technology) HRP-conjugated secondary antibody. Blots were then developed using Clarity Max chemiluminescent substrate (Bio-Rad) and protein expression was detected using a ChemiDoc MP gel imager (Bio-Rad). Densitometry was performed using Image Lab software (Bio-Rad Image Lab v 5.2). Blots were stripped and reprobed for β -Actin (#4967, 1:1000, Cell Signaling Technology) for protein normalization. Data were expressed as average volume intensity as a percentage compared to vehicle-treated E3FAD females.

2.6. Golgi impregnation and spine counting

Whole brains (n = 4–9/group) were collected for Golgi impregnation 2 weeks after the conclusion of behavioral testing. Mice

were cervically dislocated and decapitated 2 hours following DH infusion of vehicle or E_2 . The 2 hour time point was chosen because previous work from our lab and others indicates that DH infusion of E_2 increases DH and mPFC dendritic spine density at this delay (Murakami et al., 2006; Tuscher et al., 2016a). Golgi staining was performed as described previously (Frankfurt et al., 2011; Kim et al., 2019; Tuscher et al., 2016a) using the Rapid Golgi Stain Kit (FD Neuro Technologies). A cryostat was used to slice tissue into 100 μm sections, which were directly mounted onto gelatin-coated microscope slides. Mounted tissue was then stained according to the Rapid Golgi Stain Kit instructions, and coverslipped with Permount. Slides were coded so that the individual counting spines was blind to treatment, and slides were kept in the dark when not in use.

Secondary basal dendrites and tertiary apical dendrites were counted from pyramidal neurons in the dorsal hippocampal CA1 and layer II/III of the prelimbic/infralimbic mPFC under an Olympus BX51WI microscope (100x with oil) using NeuroLucida (v 11.08, MBF Bioscience). Accuracy of DH cannula placement was visually validated by examining sections containing DH tissue. Dendritic segments selected for spine counting were between 10 and 20 μm in length and 0.5–1.3 μm thick. Neurons selected for analysis were required to have well impregnated cell bodies and dendrites, and to be clearly distinguishable from adjacent cells.

Dendritic spines were identified by an experimenter scrolling through the z-plane of stained tissue in real time, and spines were reconstructed using NeuroLucida morphometric markers. Spines were classified according to 3 categories based on shape: mushroom, thin, or stubby (Harris et al., 1992; Kim et al., 2019). Mushroom spines had head diameters at least twice the size of their neck diameters, whereas thin spines had head diameters less than or equal to their neck diameters. Stubby spines had neck diameters relatively equal to the total length of the spine, with no discernable spine head. Two dendritic segments/neuron and 6 cells/region were included in the analysis. After segment tracing and spine marking, data were exported to NeuroLucida Explorer (MBF Bioscience), where a Branched Structure Analysis allowed for visualization of the number and type of spines on each traced dendritic segment. Spine density was calculated as the number of spines/10 μm dendrite.

2.7. Data analysis

All statistical analyses were conducted using GraphPad Prism 9 software (La Jolla, CA, USA). To assess within-group learning for OR and OP, one-sample *t*-tests were used to determine whether the time spent with each object during testing significantly differed from chance (15 s). Differences in memory between groups were assessed using two-way ANOVAs with treatment and genotype as between-subject variables. Similar two-way ANOVAs were used to analyze western blot and dendritic spine density data. Significant main effects were followed by Tukey's *post hoc* tests for OR, OP, western blot, and dendritic spine data. Statistical significance was set at $p < 0.05$ for all statistical tests, and trends were determined by $p < 0.10$.

3. Results

3.1. DH infusion of E_2 enhances memory in E3FAD and E3/4FAD, but not E4FAD, mice

3.1.1. Object recognition

To determine whether APOE genotype influences the ability of E_2 to facilitate memory for a previously seen object or location, ovariectomized female E3FAD, E3/4FAD, and E4FAD mice

were tested in the OR and OP tasks 24 hours or 4 hours, respectively, after receiving a post-training bilateral DH infusion of vehicle (veh; E3FAD n = 21, E3/4FAD n = 12, E4FAD n = 21) or E_2 (E3FAD n = 16, E3/4FAD n = 11, E4FAD n = 14). In OR, E_2 -treated E3FADs (Fig. 1A; $t_{(15)} = 3.406$, $p = 0.004$) and E3/4FADs (Fig. 1A; $t_{(10)} = 2.745$, $p = 0.0201$) spent significantly more time than chance (15 s) with the novel object during testing, indicating intact memory for the identity of the training objects. By contrast, E_2 -treated E4FADs, as well as all veh-treated groups, did not spend more time than chance with the novel object during testing, suggesting that 2 copies of the APOE4 allele prevented E_2 from facilitating memory. A two-way ANOVA revealed a main effect of E_2 treatment (Fig. 1A; $F_{(1, 89)} = 9.614$, $p = 0.003$), as well as a significant genotype x treatment interaction ($F_{(2, 89)} = 4.821$, $p = 0.01$) for time spent with the novel object during testing. These effects were driven by the selective benefits of E_2 on OR memory in E3FAD and E3/4FAD females only. Tukey's *post hoc* comparisons support the notion that E_2 -treated E3FAD and E3/4FADs drove the main effect of treatment, as there was a significant difference between vehicle- and E_2 -treated E3/4FADs ($p < 0.031$), and a trend for a difference between vehicle- and E_2 -treated E3FADs ($p = 0.07$).

Interestingly, time to accumulate 30 s of object exploration during testing was significantly influenced by genotype (Fig. 1C; $F_{(2, 86)} = 11.93$, $p < 0.0001$), such that E4FADs took significantly less time to reach 30 seconds of object exploration relative to other groups. E4FADs treated with E_2 were particularly fast to complete object exploration, as demonstrated by *post hoc* comparisons to other E_2 -treated groups (Fig. 1C; $p = 0.002$ vs. E_2 -treated E3FADs, and $p = 0.0004$ vs. E_2 -treated E3/4FADs). Given the significant genotype effect during testing, we also examined time to accumulate 30 seconds of object exploration during training, but found no significant effects (Veh-treated E3FADs, $M = 541.35$, SEM = 56.359; E_2 -treated E3FADs, $M = 537.957$, SEM = 48.977; Veh-treated E3/4FADs, $M = 447.892$, SEM = 65.291; E_2 -treated E3/4FADs, $M = 538.325$, SEM = 70.724; Veh-treated E4FADs, $M = 557.613$, SEM = 71.566; E_2 -treated E4FADs, $M = 382.463$, SEM = 57.742). This pattern of data suggests that E_2 -treated E4FADs were not generally hyperactive, but rather just quick to explore objects during testing.

3.1.2. Object placement

The OP task yielded similar results to OR in that E_2 treatment selectively improved memory in E3FAD and E3/4FAD females. E_2 -treated E3FAD (Fig. 1B; $t_{(12)} = 3.741$, $p = 0.003$) and E3/4FAD (Fig. 1B; $t_{(10)} = 5.644$, $p = 0.0002$) females spent significantly more time than chance with the moved object during testing, indicating intact memory for the location of the training objects. As in OR, E_2 -treated E4FAD and veh-treated mice of all genotypes did not spend more time than chance with the moved object during testing, again suggesting that APOE4 homozygotes were unresponsive to the mnemonic benefits of E_2 . As with OR, the main effect of treatment was significant (Fig. 1B; $F_{(1, 88)} = 8.183$, $p = 0.005$), as was the genotype x treatment interaction (Fig. 1B; $F_{(2, 88)} = 3.591$, $p = 0.032$), reflecting the selective benefit of E_2 for E3FAD and E3/4FAD females. However, unlike OR, the main effect of genotype was also significant (Fig. 1B; $F_{(2, 88)} = 10.72$, $p < 0.0001$), such that E3FAD and E3/4FADs exhibited better memory for object location than E4FADs. In support of these conclusions, Tukey's *post hoc* comparisons indicated that E_2 -treated E4FADs spent significantly less time with the moved object than E_2 -treated E3FADs (Fig. 1B; $p = 0.002$) and E3/4FADs (Fig. 1B; $p = 0.0009$).

Similar to OR, time to accumulate 30 seconds of object exploration during testing was significantly modulated by genotype (Fig. 1D; $F_{(2, 89)} = 8.765$, $p = 0.0003$), such that E4FADs were faster to accumulate 30 seconds of object exploration relative to




Fig. 1. E₂ facilitates object recognition and spatial memory formation in E3FAD and E3/4FAD, but not E4FAD, females. (A, B) E3FAD and E3/4FAD females treated with E₂ spent significantly more time than chance (dashed line at 15 sec; $^{\wedge} p < 0.05$, $^{\wedge\wedge} p < 0.01$, $^{\wedge\wedge\wedge} p < 0.001$) with the novel (A) and moved (B) objects during testing. In contrast, E₂-treated E4FAD females did not, nor did any groups treated with vehicle (veh). The genotype-dependent effect of E₂ on memory was also reflected in between-subjects' analyses by a main effect of treatment ($^{\$\$} p < 0.01$) and treatment \times genotype interaction ($^{*\wedge} p < 0.05$). In OP, the main effect of genotype ($^{&&&} p < 0.0001$) indicated that E3FADs and E3/4FADs spent more time with the moved object than E4FADs. Tukey's post hoc comparisons revealed significant differences between veh- and E₂-treated E3/4FADs in OR (A; $*p < 0.05$), and that E₂-treated E3FADs ($^{**} p < 0.01$) and E3/4FADs ($^{***} p < 0.001$) spent more time with the moved object in OP than E₂-treated E4FADs. (C, D) The main effect of genotype was significant for both OR (C; $^{&&&} p < 0.0001$) and OP (D; $^{&&} p < 0.001$). E4FADs took less time to accumulate 30 sec of exploration in both tasks. Tukey's post hoc comparisons indicated that E₂-treated E4FADs were faster than E₂-treated E3/4FADs and E3FADs in OR (C; $^{**} p < 0.01$), and than E₂-treated E3FADs in OP (D; $^{**} p < 0.01$). Bars represent mean \pm standard error of the mean (SEM).

other groups. Exploration time in E₂-treated E4FADs was significantly lower than that of E₂-treated E3FADs (Fig. 1D; $p = 0.007$). We again examined time to accumulate 30 seconds of object exploration during training, but found no significant effects (Veh-treated E3FADs, $M = 495.968$, SEM = 59.455; E₂-treated E3FADs, $M = 400.2$, SEM = 56.820; Veh-treated E3/4FADs, $M = 431.2$, SEM = 77.794; E₂-treated E3/4FADs, $M = 399.033$, SEM = 34.655; Veh-treated E4FADs, $M = 430.119$, SEM = 67.464; E₂-treated E4FADs, $M = 326.927$, SEM = 36.685).

Combined, data from both OR and OP tasks suggest that E₂ supports memory consolidation in E3FAD and E3/4FAD, but not E4FAD, females.

3.1.3. ER α is elevated in the DH of E4FAD mice

Because we previously found that APOE4 increased ER α expression in the DH of gonadally-intact EFAD mice of both sexes (Taxier et al., 2022a), we examined effects of DH vehicle or E₂ infusion on levels of this protein and 2 other estrogen receptors, ER β and G-protein-coupled estrogen receptor (GPER). Two-way ANOVA revealed a significant main effect of genotype, but not treatment, nor an interaction, for levels of ER α (Fig. 2A,B; $F_{(2,48)} = 6.956$,

$p = 0.002$), driven by elevated ER α in E₂-treated E4FADs relative to veh-treated E3FADs ($p = 0.019$ veh-treated E3FADs vs. E₂-treated E4FADs, $p = 0.052$ veh-treated E3FADs vs. veh-treated E4FADs). There were no significant effects of treatment or genotype in DH protein expression of either ER β or GPER (Fig. 2A,C,D).

3.1.4. ERK and JNK are not rapidly phosphorylated by E₂ in EFAD mice

In wildtype ovariectomized C57BL/6 mice, the memory-enhancing effects of E₂ are facilitated by ER α and ER β signaling via phosphorylation of extracellular signal-regulated kinase (ERK) (Boulware et al., 2013). Although activation of GPER also facilitates memory in ovariectomized mice, it does so independently from ERK and from DH E₂ infusion via c-jun N-terminal kinase (JNK) activation (Kim et al., 2016). Because the cell-signaling events underlying E₂-facilitated memory are uncharacterized in EFADs, we examined whether E₂ facilitates memory via activation of ERK or JNK signaling in this model. Surprisingly, there were no significant effects of E₂ or genotype on ERK or JNK phosphorylation in the DH (Table 1).




Fig. 2. ER α levels in the DH were increased in E4FAD females. (A) Representative western blot images. (B) ER α was affected by APOE genotype ($^{&&}p < 0.01$ = main effect of genotype) such that levels were highest in E4FAD females. ER α levels were significantly higher in E $_2$ -treated E4FAD mice than in veh-treated E3FAD mice (* $p < 0.05$). (C,D) Levels of ER β (C) and GPER (D) were unaffected by treatment or genotype.

Table 1
Levels of phosphorylated ERK and JNK were unaffected by E $_2$ treatment or genotype

Protein*	E3FAD		E3/4FAD		E4FAD	
	Veh	E $_2$	Veh	E $_2$	Veh	E $_2$
p42 ERK	100.0 ± 10.56	93.21 ± 10.05	109.39 ± 20.84	112.56 ± 16.44	109.45 ± 15.76	110.17 ± 14.46
p44 ERK	100.0 ± 9.32	119.51 ± 8.41	97.37 ± 12.20	124.85 ± 9.37	133.63 ± 18.73	111.44 ± 9.96
p46 JNK	100.0 ± 2.05	89.52 ± 3.88	93.15 ± 6.28	94.97 ± 4.50	102.83 ± 5.25	94.02 ± 4.60
p54 JNK	100.0 ± 5.86	89.70 ± 7.51	96.39 ± 8.67	90.60 ± 7.57	97.68 ± 7.94	8.35 ± 8.05

* All proteins were normalized to the vehicle-treated E3FAD group, whose values were set to 100. Values represent mean % immunoreactivity (± SEM) for the target phosphorylated protein divided by unphosphorylated total protein.

3.2. DH and mPFC spine densities are affected by APOE genotype and E $_2$ treatment

We next examined apical and basal dendritic spine density on pyramidal neurons in hippocampal area CA1 and in the prelimbic/infralimbic area of the mPFC to assess whether APOE genotype influenced the ability of E $_2$ to increase dendritic spine density 2 hours following DH infusion. Representative images of vehicle- and E $_2$ -treated spines for each genotype are shown in Figs. 3A, 4A, and 5A, whereas representative images of mushroom, stubby, and thin spines are shown in Fig. 3B. There were no effects of genotype or E $_2$ treatment, nor an interaction, on mPFC apical spines (Table 2). Treatment and genotype effects on CA1 apical, and CA1 and mPFC basal spines, are detailed below.

3.2.1. CA1

APOE4 homozygosity blocked E $_2$'s effects on CA1 total apical spine density, as suggested by the fact that E $_2$ significantly increased CA1 total apical spine density in E3FADs and E3/4FADs, but not E4FADs. Significant main effects of genotype (Fig. 3C; $F_{(2,29)} = 6.746$, $p = 0.0039$) and E $_2$ treatment (Fig. 3C; $F_{(1,29)} = 30.25$, $p < 0.0001$) were observed, as well as a genotype

x treatment interaction (Fig. 3C; $F_{(2,29)} = 4.536$, $p = 0.0193$). Tukey post hoc tests revealed that total apical spine density was significantly higher in E3FAD ($p = 0.0032$) and E3/4FAD ($p = 0.0033$) mice treated with E $_2$ than with vehicle, whereas there were no effects of E $_2$ in E4FAD mice ($p = 0.932$). Moreover, apical spine density was significantly lower in E $_2$ -treated E4FAD mice than in E $_2$ -treated E3FAD ($p = 0.0451$) and E3/4FAD mice ($p = 0.002$). Further analysis by spine subtype revealed no significant effects of genotype or treatment, nor interactions, on CA1 apical thin or mushroom spines (Table 2). However, the main effects of genotype (Fig. 3D; $F_{(2,31)} = 4.261$, $p = 0.0002$) and treatment (Fig. 3D; $F_{(1,31)} = 9.712$, $p = 0.012$) were significant for CA1 apical stubby spines. Here, the only significant between-group differences were between the E3FAD and E4FAD vehicle groups, such that veh-treated E3FAD mice had more stubby apical spines than veh-treated E4FAD mice ($p = 0.006$).

CA1 basal spine density was influenced by genotype only. The main effect of genotype was significant (Fig. 4B; $F_{(2,31)} = 7.774$, $p = 0.002$), but the treatment effect and interaction were not. This effect was driven largely by elevated total spine density in E $_2$ -treated E3FAD mice, who had significantly higher basal spine density than E $_2$ -treated E4FADs (Fig. 4B; $p = 0.002$). Further anal-




Fig. 3. E₂ increased CA1 apical spine density in E3FAD and E3/4FAD, but not E4FAD, females. (A) Representative images of CA1 apical dendritic spine segments at 100X magnification. (B) Representative images of stubby (top left), thin (top right), and mushroom (bottom) spines. Arrows indicate individual examples of each spine class. (C) Total CA1 apical dendritic spine density was increased by DH E₂ infusion ($$$$$p < 0.01$ = main effect of treatment) and reduced by 2 copies of APOE4 ($\&\&p < 0.05$ = main effect of genotype; $\%p < 0.05$ = interaction). In both E3FAD and E3/4FAD females, E₂ treatment increased total apical spine density relative to veh ($**p < 0.01$), but no benefit of E₂ was observed in E4FAD mice. Moreover, apical spine density was significantly lower in E₂-treated E4FAD mice than in E₂-treated E3FAD ($*p < 0.05$) and E3/4FAD ($**p < 0.01$) mice. (C) CA1 apical stubby spines were increased by E₂ ($\$p < 0.05$ = main effect of treatment) and reduced by 2 copies of APOE4 ($\&\&p < 0.001$ = main effect of genotype). No differences were observed between veh and E₂ within any genotype, but veh-treated E3FADs had significantly more stubby spines than veh-treated E4FADs ($**p < 0.01$).




Fig. 4. CA1 basal spine density was reduced by APOE4. (A) Representative images of CA1 basal dendritic spine segments at 100X magnification. (B-E) Total CA1 basal spine density (B), stubby spine density (C), and thin spine density (D) were decreased in E4FADs regardless of treatment ($^8p < 0.05$, $\&\&p < 0.01$ = main effect of genotype). The treatment x genotype interaction was significant for mushroom spines (E; $\%p < 0.05$). E₂-treated E3FAD females had more total spines, thin spines, and mushroom spines than E₂-treated E4FAD females ($*p < 0.05$, $**p < 0.01$).




Fig. 5. mPFC basal spine density was reduced by *APOE4*. (A) Representative images of mPFC basal dendritic spine segments at 100X magnification. (A) *APOE4* genotype reduced mPFC basal spines, as suggested by a modest main effect of genotype for total spines (B; $^{\&}p = 0.055$) and a significant main effect in thin spines (C; $^{\&}p < 0.05$). No between-group differences were observed.

Table 2
CA1 and mPFC spine density measures unaffected by E_2 treatment or genotype

Brain region	Spine type	E3FAD		E3/4FAD		E4FAD	
		Veh	E_2	Veh	E_2	Veh	E_2
mPFC	Apical thin	3.69 ± 0.18	4.51 ± 0.29	4.06 ± 0.28	4.02 ± 0.11	4.26 ± 0.27	4.32 ± 0.21
	Apical mushroom	3.05 ± 0.11	3.06 ± 0.25	3.30 ± 0.23	3.71 ± 0.30	3.52 ± 0.28	3.20 ± 0.11
	Apical total	11.6 ± 0.88	10.35 ± 0.46	11.43 ± 0.95	10.03 ± 0.72	10.19 ± 0.85	9.84 ± 0.48
	Apical thin	2.74 ± 0.24	2.27 ± 0.06	2.41 ± 0.26	2.24 ± 0.10	2.75 ± 0.28	2.51 ± 0.14
	Apical stubby	5.32 ± 0.48	4.77 ± 0.32	5.42 ± 0.43	4.95 ± 0.59	4.59 ± 0.55	4.46 ± 0.38
	Basal stubby	7.36 ± 0.46	6.63 ± 0.28	6.80 ± 0.26	6.95 ± 0.57	7.82 ± 0.38	6.79 ± 0.24
	Basal mushroom	2.84 ± 0.20	2.96 ± 0.23	2.42 ± 0.32	2.73 ± 0.50	2.32 ± 0.27	2.61 ± 0.29

Values represent mean spines/10 $\mu\text{m} \pm \text{SEM}$

ysis by spine subtype revealed that the main effect of genotype on total CA1 basal spine density was driven by significant effects of genotype on CA1 basal stubby spines (Fig. 4C; $F_{(2,31)} = 4.49$, $p = 0.019$) and thin spines (Fig. 4D; $F_{(2,31)} = 7.887$, $p = 0.002$). E_2 -treated E3FADs exhibited elevated thin basal spines relative to E_2 -treated E4FADs (Fig. 4D; $p = 0.013$). The genotype x treatment interaction was significant for CA1 basal mushroom spines (Fig. 4E; $F_{(2,30)} = 5.151$, $p = 0.012$), such that mushroom spine density was elevated in E_2 -treated E3FADs relative to E_2 -treated E4FADs (Fig. 4E; $p = 0.012$).

3.2.2. mPFC

Neither genotype nor E_2 treatment affected mPFC apical spine density in terms of total spines or spine subtypes (Table 2). However, there were modest effects of genotype on mPFC basal spines, such that E4FADs again had the lowest spine density. For total basal spines, a two-way ANOVA revealed a trend towards a significant main effect of genotype (Fig. 5B; $F_{(2,30)} = 3.178$, $p = 0.055$), which appeared to be driven by a significant reduction among E4FAD mice in thin spines (Fig. 5C; $F_{(2,30)} = 5.212$, $p = 0.011$). Although no post hoc comparisons were significant, E3FADs gen-

erally had elevated mPFC basal thin spines relative to EFADs of other genotypes, regardless of treatment. There were no significant genotype, treatment, nor genotype x treatment effects on levels of mPFC basal stubby or mushroom spines (Table 2).

4. Discussion

Because *APOE4* status and female sex are unmodifiable risk factors that act synergistically to increase AD risk (Altmann et al., 2014), evaluating interventional approaches that might benefit female *APOE4* carriers is a crucial step towards providing individualized treatment to AD patients. Here, we investigated whether *APOE* genotype interacts with the ability of E_2 to promote memory consolidation, modulate estrogen receptor expression, and increase CA1 or mPFC dendritic spine density in an EFAD mouse model of AD designed to model *APOE*-associated disease risk against a background of AD-like pathology. We hypothesized that E_2 would have the most beneficial effects on memory consolidation and spine density in ovariectomized E3FAD mice, and that E_2 may exert lesser effects in E3/4FAD mice due to the presence of one *APOE4* allele. Based on data from women showing no protective effect of

estrogen therapy on cognitive decline in *APOE4* carriers (Yaffe et al., 2000), we expected that E₂ would have little to no influence on memory or spine density in E4FAD females and intermediate effects in E3/4FAD mice due to the presence of one *APOE4* allele. We found that DH infusion of E₂ facilitated object memory consolidation in both E3FAD and E3/4FAD mice, but not E4FAD mice, indicating that 2 copies of the *APOE4* allele blocks the memory-enhancing effects of E₂. The effects of E₂ in E3FAD and E3/4FAD females were associated with an increase in total CA1 apical spine density. Independent of E₂ treatment, *APOE4* genotype alone reduced CA1 and mPFC spine density and increased levels of ER α in the DH.

The beneficial effects of acute DH E₂ infusion on object recognition and spatial memory consolidation in 6-month-old ovariectomized E3FAD and E4FAD mice are consistent with an extensive literature from our lab and others showing similar benefits of DH E₂ infusion in 2–3-month-old wild-type ovariectomized C57BL/6 mice (Fernandez et al., 2008; Fortress et al., 2013; Gross et al., 2021; Packard and Teather, 1997; Taxier et al., 2019). A primary difference is that the delays between training and testing used here (24 hours for OR and 4 hours for OP) were shorter than those at which E₂ benefits wild-types (48 hours for OR and 24 hours for OP), but this difference is to be expected given the significant AD-like pathology present in the brains of EFAD mice (Tai et al., 2017; Youmans et al., 2012). Although others have shown a potential benefit of chronic E₂ treatment for decreasing amyloid accumulation in ovariectomized EFAD mice (Kunzler et al., 2014) and in ovariectomized 3xTg mice (Carroll et al., 2007), our data are the first to demonstrate that a single acute dose of E₂ can facilitate memory consolidation in E3FAD and E3/4FAD, but not E4FAD mice. Previous work demonstrated that chronic E₂ treatment initiated in 3-month-old EFADs can significantly reduce amyloid burden (Kunzler et al., 2014). We find it particularly striking that E3FAD and E3/4FAD mice remain responsive to a single acute dose of E₂ until at least 6 months of age, at which point there is already substantial AD-like pathology evident (Tai et al., 2017). Although recent clinical studies suggest that estrogen therapy is not efficacious for women with AD, these findings are clouded by differences in treatment regimen, age of experimental subjects and menopausal status, and lack of stratification by *APOE* status in some instances. Thus, whether *APOE3+* women with AD who express 0 or 1 copies of *APOE4* can benefit cognitively from E₂ treatment after substantial disease progression remains a question for future study.

Although there was a significant interaction between genotype and treatment in both the OR and OP tasks, the pro-cognitive effects of E₂ were more apparent for OP, suggesting that spatial memory may be more sensitive to E₂ in ovariectomized E3FAD and E3/4FAD mice than recognition memory. Alternatively, the longer 24-hour delay between training and testing for OR may have been more demanding on memory function than the shorter 4-hour delay for OP. Interestingly, E4FAD mice took less time than other genotypes to accumulate 30 seconds of object exploration during testing, but not during training, which is an unusual finding in our extensive experience using these tasks to examine estrogenic regulation of memory. One possible explanation is that E4FAD mice noticed a change during testing that caused an increase in exploratory behavior, but without the ability to discriminate between old and new objects and locations, this increase did not result in a preference for novelty. Our previous work with gonadally-intact female EFAD mice indicated that during object memory habituation trials, E4FAD females explored the outer third of the open field box more than E3FAD females (Taxier et al., 2022b). Our objects are located in this peripheral zone, and so a possible training-induced increase in exploration in this zone could have caused this group to contact and explore the objects faster than other groups that

were exploring the entirety of the testing arena. Future studies will need to further address this possibility.

Importantly, our results also suggest that individuals carrying a single copy of the *APOE4* allele may still benefit from the cognitive benefits of E₂ treatment, whereas individuals with 2 copies would not. This finding is somewhat consistent with data from human patients, from which the conclusion is sometimes reached that estrogens are ineffectual or in some instances harmful for cognition in *APOE4* carriers (Burkhardt et al., 2004; Kang and Grodstein, 2012). Others have reported neuroprotective effects and cognitive benefits of estrogens in human *APOE4* carriers, especially when treatment is initiated within a critical window (Jacobs et al., 2013; Mahoney et al., 2020). Because individuals carrying 2 *APOE4* alleles are relatively rare in human populations (Heffernan et al., 2016), therefore making sufficiently-powered sample size a challenge, studies with human patients do not often differentiate between women carriers of one or 2 *APOE4* alleles (but see Mahoney et al., 2020). Data from the present study suggests that this distinction is important because heterozygous E3/4FAD female mice responded to E₂, whereas homozygous E4FADs did not. More work remains to determine what factors mediate the neuroprotective properties of E₂ in carriers of one or both copies of an *APOE4* allele.

Surprisingly, the memory-enhancing effects of E₂ in the present study occurred in the absence of rapid p42 ERK phosphorylation in the DH, which has been documented many times in young adult wildtype C57BL/6 mice and is necessary in these mice for E₂ to enhance memory consolidation in the object recognition and object placement tasks (e.g., Boulware et al., 2013; Fernandez et al., 2008; Fortress et al., 2013). Therefore, data from the current study suggest that the benefits of E₂ in ovariectomized E3FAD and E3/4FAD mice occur either via a different time course of ERK activation or an as-of-yet uncharacterized estrogen-sensitive signaling event. The absence of rapid JNK phosphorylation in the present study also suggests GPER-independent signaling, given that the effects of GPER activation on object recognition and object placement memory consolidation in young ovariectomized wildtype mice are dependent upon activation of this kinase (Kim et al., 2016, 2019). It is possible that pathological changes in the DH of these mice (e.g., Youmans et al., 2012) altered the timing of kinase activation relative to wild types, so collecting tissue from E₂-infused EFADs at extended time points would allow for a more complete picture of E₂-related cell signaling across genotypes.

Our previous study of gonadally-intact male and female E3FAD and E4FAD mice revealed that E4FADs of both sexes had significantly higher levels of ER α in the DH (Taxier et al., 2022a). The present study is consistent with this finding and extends it to ovariectomized E4FAD mice. Our data thus far suggest that the increase in ER α among E4FAD mice is associated with impaired memory consolidation (Taxier et al., 2022a) and an inability of E₂ to enhance memory consolidation (present study). Consistent with these data, others have shown that ER α is elevated in the hippocampus of AD patients, particularly *APOE4* carriers (Ishunina and Swaab, 2003). It is unclear why this is so. It could be that ER α is increased in response to changes in levels of other ERs. However, levels of ER β and GPER were unaffected by genotype or treatment, so this is unlikely. The increase in ER α could come in response to reduced levels of E₂ synthesized within the hippocampus. Levels of E₂ and mRNA levels of the E₂-synthesizing enzyme aromatase were lower in frontal cortices and cerebella of postmortem brains from female AD patients than in non-AD controls (Yue et al., 2005), supporting the possibility that ER α levels may increase in an attempt to compensate for lower levels of brain E₂. However, it is unknown if or how levels of E₂ or aromatase are altered by genotype

in EFAD mice. Thus, the relationship between *APOE4* homozygosity and ER α levels warrants future study, especially in light of previous findings from rat hippocampus that *APOE* mRNA and *APOE* protein expression are upregulated 24 hours following ER α activation, either with E₂ or an ER α agonist (Wang et al., 2006). Future work should examine whether hippocampal *APOE* expression is altered following E₂ treatment in E3FAD, E3/4FAD, or E4FADs. With respect to the lack of E₂ effects on ER levels in the present study, it should be noted that 5 minutes is a very rapid timepoint at which to see effects on receptor expression of a single E₂ treatment, so future studies should examine longer time points and/or longer durations of E₂ treatment. Better understanding the extent to which levels of the 3 ERs are altered by genotype and in response to E₂ treatment across genotypes could provide valuable information as to whether selective estrogen receptor modulators could be potentially useful therapeutics for preventing or treating cognitive decline in *APOE4*-positive women.

The effects of E₂ on CA1 apical dendritic spine density were remarkably consistent with its beneficial effects on memory. In line with its memory-enhancing effects in ovariectomized E3FAD and E3/4FAD females, E₂ increased CA1 total apical spine density in E3FAD and E3/4FAD, but not E4FAD, females 2 hours after infusion. This effect is consistent with previous findings from our lab showing that CA1 total apical spine density is increased 2 hours after E₂ infusion in wild-type ovariectomized mice (Tuscher et al., 2016a). Here, we also showed that E₂ modestly increased CA1 apical stubby spine density in EFADs, although levels of apical stubby spines were similar in E3FADs regardless of treatment. Although thin spines are typically considered more likely than stubby spines to transition to other spine types, previous work in another AD model suggests that stubby spines can similarly take on this dynamic role in pathological conditions (Spires-Jones et al., 2007). Therefore, an increase in apical stubby spines in CA1 may reflect ongoing spine remodeling or spinogenesis stimulated by E₂. Interestingly, E₂ also increases CA1 and mPFC basal spines 2 hours post intrahippocampal E₂-infusion in wild-type ovariectomized mice (Tuscher et al., 2016a). However, we did not see a corresponding increase in CA1 and mPFC basal spines following E₂ infusion in E3FADs or E3/4FADs, which may be reflective of reduced interactions or connectivity between mPFC and CA1 in E₂-treated EFADs relative to wild-type mice. Nevertheless, our analyses suggest that E₂ may exert its memory-enhancing effects in E3FAD and E3/4FAD mice via increased CA1 apical spine density.

E₂ also had a striking effect on CA1 basal mushroom spines, for which the genotype \times treatment interaction was significant. This finding, in which E₂-treated ovariectomized E3FADs had the highest number of CA1 basal mushroom spines, whereas E₂-treated E4FADs had the lowest, represents a key measure by which E₂ benefits *APOE3* carriers, while harming *APOE4* carriers. Because mushroom spines are thought to be mature spines (Bourne and Harris, 2007; Hayashi and Majewska, 2005), the beneficial effects of E₂ on memory in E3FADs, and its inefficacy in E4FADs, may be linked to the respective E₂-induced increase of CA1 basal mushroom spines in E3FADs, and E₂-induced decrease of this same spine subtype in E4FADs. That is, in addition to effects on CA1 apical spines, E₂ may facilitate memory in E3FADs by increasing the density of mature spines on CA1 basal dendrites and impair memory in E4FADs by reducing basal mature spine density. However, this possibility is limited by the fact that the same effect of E₂ on CA1 basal mushroom spines was not present in E3/4FADs at the single time point we examined. Recent work suggests that dendritic spine loss does not occur at random in AD patients; rather, spines are lost in clusters depending on the presence or absence of tau pathology in examined cells (Mijalkov et al., 2021). Therefore, because cells and dendritic segments in the present

study were semi-randomly chosen for analyses based on stain penetrance, our analyses are limited by the fact that we do not know for certain whether spines were counted from neurons in which AD-like pathology was present. The use of advanced methods such as immuno-EM in tissue collected from EFADs could provide ultrastructural information about the presence or absence of pathological hallmarks of AD in specific cellular compartments (Takahashi et al., 2010).

Genotype modulated spine density in a manner largely consistent with our hypotheses, in that genotype effects on CA1 apical, CA1 basal, and mPFC basal spines were seemingly driven by increased dendritic spine density in E3FADs, or decreased dendritic spine density in E4FADs, relative to other groups, which may have contributed to their impaired memory consolidation. The main effect of genotype on basilar thin spine density in CA1 is of particular interest because *post hoc* comparisons indicated that E₂-treated E3FADs had the highest density of this subtype. Our data are consistent with previous work showing that treatment with E₂ facilitated cognitive function in aged rhesus monkeys, an effect dependent upon increased thin spine density (Hao et al., 2006). Furthermore, our data fit with the notion that thin spines are thought to be “learning spines,” or transient protrusions that often evolve into more mature, mushroom “memory” spines (Bourne and Harris, 2007). CA1 basal and proximal apical dendrites receive projections from CA3 to support memory processes (Li et al., 1994; Spruston, 2008; Whitlock et al., 2006), and E₂ treatment evokes an increase in synaptic transmission at CA3-CA1 synapses (Kumar et al., 2015; Smejkalova and Woolley, 2010; Smith et al., 2009). Therefore, higher density of CA1 total apical spines in E₂-treated E3FADs and E3/4FADs, as well as higher density of CA1 basilar thin spines in E₂-treated E3FADs, could be linked to improved memory processes. The lack of genotype or treatment effects on mPFC apical spine density (Table 2), and modest genotype effects on mPFC basal spine density indicate that the memory-enhancing effects of E₂ in ovariectomized E3FADs and E3/4FADs are not mediated through changes in mPFC spine density. However, the genotype effects on mPFC total and thin basal spines suggests that E3FAD females have somewhat higher basal spine density relative to E4FAD females, which is consistent with our previous findings in gonadally-intact E3FAD and E4FAD mice (Taxier et al., 2022a).

In sum, intrahippocampal E₂ infusion facilitated memory consolidation and increased CA1 apical spine density in ovariectomized E3FAD and E3/4FAD mice, suggesting that mice with AD-like pathology bearing 0 or 1 copy of *APOE4* remain responsive to acute E₂ treatment. Notably, 2 copies of *APOE4* had a deleterious effect on the ability of E₂ to enhance memory and increase dendritic spine density, and also increased levels of ER α in a manner possibly linked to impaired memory consolidation. In addition, the fact that E₂ increased CA1 basal mushroom spine density in E3FADs and reduced these spines in E4FADs may account for the respective ability and inability of E₂ to enhance memory in E3FADs and E3/4FADs versus E4FADs. These data support the notion that E₂ treatment can potentially benefit individuals who carry 0 or 1 copy of the *APOE4* allele, but not those with 2 copies. Future experiments should examine the kinetics of both protein expression and dendritic spine density following E₂ treatment at multiple time points to fully elucidate the molecular correlates of the mnemonic benefits of E₂ in EFAD mice. Much more work remains to fully characterize the interactive effects between E₂ and *APOE4* genotype and to determine how *APOE4* impedes the actions of E₂. This information may provide new important new insights about the impact of E₂ loss on AD risk in women and better define subpopulations of women who could benefit cognitively from estrogen therapy.

Disclosure Statement

K.M.F. is a co-founder and the Chief Scientific Officer of Estringenix Therapeutics, Inc. The other authors have no actual or potential conflicts of interest to declare.

Acknowledgments

We gratefully acknowledge the primary support of an inaugural Sex and Gender in Alzheimer's grant ([SAGA-17- 419092](#)) from the Alzheimer's Association for this work. The Frick lab was also supported by [R01MH107886](#) to KMF, [2R15GM118304-02](#) to KMF, and [1F31MH118822-01A1](#) to LRT, the UWM College of Letters & Science, and the UWM Office of Undergraduate Research. Additional funding of the LaDu lab was provided by [R01AG056472](#), [R01AG057008](#), [UH2/3NS100127](#), [R56AG058655](#), philanthropic support from Lou and Christine Friedrich, and UIC institutional funds. We also thank Dr. James R. Moyer, Jr. for use of his Olympus BX51WI microscope and NeuroLucida software.

CRediT authorship contribution statement

Lisa R. Taxier: Conceptualization, Methodology, Validation, Formal analysis, Investigation, Writing – original draft, Writing – review & editing, Visualization, Project administration. **Sarah M. Philippi:** Formal analysis, Investigation, Writing – original draft, Writing – review & editing. **Aaron W. Fleischer:** Investigation, Writing – review & editing. **Jason M. York:** Resources, Writing – review & editing. **Mary Jo LaDu:** Conceptualization, Methodology, Resources, Writing – review & editing, Visualization, Funding acquisition. **Karyn M. Frick:** Conceptualization, Methodology, Validation, Writing – original draft, Writing – review & editing, Visualization, Supervision, Project administration, Funding acquisition.

References

- Altmann, A., Tian, L., Henderson, V.W., Greicius, M.D., 2014. Sex modifies the APOE-related risk of developing Alzheimer disease. *Ann. Neurol.* 75, 563–573. doi: [10.1002/ana.24135](#).
- Baldereschi, M., Di Carlo, A., Lepore, V., Bracco, L., Maggi, S., Grigoletto, F., Scarlato, G., Amaducci, L., 1998. Estrogen-replacement therapy and Alzheimer's disease in the Italian longitudinal study on aging. *Neurology* 50, 996–1002. doi: [10.1212/wnl.50.4.996](#).
- Bertram, L., 2009. Alzheimer's disease genetics current status and future perspectives. *Int. Rev. Neurobiol.* 84, 167–184. doi: [10.1016/S0074-7742\(09\)00409-7](#).
- Boulware, M.I., Heisler, J.D., Frick, K.M., 2013. The memory-enhancing effects of hippocampal estrogen receptor activation involve metabotropic glutamate receptor signaling. *J. Neurosci.* 33, 15184–15194. doi: [10.1523/JNEUROSCI.1716-13.2013](#).
- Bourne, J., Harris, K.M., 2007. Do thin spines learn to be mushroom spines that remember? *Curr. Opin. Neurobiol.* 17, 381–386. doi: [10.1016/j.conb.2007.04.009](#).
- Bretsky, P.M., Buckwalter, J.G., Seeman, T.E., Miller, C.A., Poirier, J., Schellenberg, G.D., Finch, C.E., Henderson, V.W., 1999. Evidence for an interaction between apolipoprotein E genotype, gender, and Alzheimer disease. *Alzheimer Dis. Assoc. Disord.* 13, 216–221. doi: [10.1097/00002093-199910000-00007](#).
- Burkhardt, M.S., Foster, J.K., Laws, S.M., Baker, L.D., Craft, S., Gandy, S.E., Stuckey, B.G.A., Clarnette, R., Nolan, D., Hewson-Bower, B., Martins, R.N., 2004. Oestrogen replacement therapy may improve memory functioning in the absence of APOE ε4. *J. Alzheimers Dis.* 6, 221–228. doi: [10.3233/JAD-2004-6302](#).
- Carroll, J.C., Rosario, E.R., Chang, L., Stanczyk, F.Z., Oddo, S., LaFerla, F.M., Pike, C.J., 2007. Progesterone and estrogen regulate Alzheimer-like neuropathology in female 3xTg-AD mice. *J. Neurosci.* 27, 13357–13365. doi: [10.1523/JNEUROSCI.2718-07.2007](#).
- DeKosky, S.T., Scheff, S.W., 1990. Synapse loss in frontal cortex biopsies in Alzheimer's disease: correlation with cognitive severity. *Ann. Neurol.* 27, 457–464. doi: [10.1002/ana.410270502](#).
- Dumanis, S.B., Tesoriero, J.A., Babus, L.W., Nguyen, M.T., Trotter, J.H., LaDu, M.J., Weeber, E.J., Turner, R.S., Xu, B., Rebeck, G.W., Hoe, H.-S., 2009. ApoE4 decreases spine density and dendritic complexity in cortical neurons *in vivo*. *J. Neurosci.* 29, 15317–15322. doi: [10.1523/JNEUROSCI.4026-09.2009](#).
- Fernandez, S.M., Lewis, M.C., Pechenino, A.S., Harburger, L.L., Orr, P.T., Gresack, J.E., Schafe, G.E., Frick, K.M., 2008. Estradiol-induced enhancement of object memory consolidation involves hippocampal extracellular signal-regulated kinase activation and membrane-bound estrogen receptors. *J. Neurosci.* 28, 8660–8667. doi: [10.1523/JNEUROSCI.1968-08.2008](#).
- Fortress, A.M., Fan, L., Orr, P.T., Zhao, Z., Frick, K.M., 2013. Estradiol-induced object recognition memory consolidation is dependent on activation of mTOR signaling in the dorsal hippocampus. *Learn. Mem.* 20, 147–155. doi: [10.1101/lm.026732.112](#).
- Frankfurt, M., Salas-Ramirez, K., Friedman, E., Luine, V., 2011. Cocaine alters dendritic spine density in cortical and subcortical brain regions of the postpartum and virgin female rat. *Synapse* 65, 955–961. doi: [10.1002/syn.20918](#).
- Gealon, C.E., Dowling, N.M., Wharton, W., Manson, J.E., Miller, V.M., Atwood, C.S., Brinton, E.A., Cedars, M.I., Lobo, R.A., Merriam, G.R., Neal-Perry, G., Santoro, N.F., Taylor, H.S., Black, D.M., Budoff, M.J., Hodis, H.N., Naftolin, F., Harman, S.M., Asthana, S., 2015. Effects of hormone therapy on cognition and mood in recently postmenopausal women: findings from the randomized, controlled KEEPS-cognitive and affective study. *PLoS Med.* 12, e1001833. doi: [10.1371/journal.pmed.1001833](#), discussion e1001833.
- Gross, K.S., Alf, R.L., Polzin, T.R., Frick, K.M., 2021. 17β-estradiol activation of dorsal hippocampal TrkB is independent of increased mature BDNF expression and is required for enhanced memory consolidation in female mice. *Psychoneuroendocrinology* 125, 105110. doi: [10.1016/j.psyneuen.2020.105110](#).
- Hao, J., Rapp, P.R., Leffler, A.E., Leffler, S.R., Janssen, W.G.M., Lou, W., McKay, H., Roberts, J.A., Wearne, S.L., Hof, P.R., Morrison, J.H., 2006. Estrogen alters spine number and morphology in prefrontal cortex of aged female rhesus monkeys. *J. Neurosci.* 26, 2571–2578. doi: [10.1523/JNEUROSCI.3440-05.2006](#).
- Harris, K.M., Jensen, F.E., Tsao, B., 1992. Three-dimensional structure of dendritic spines and synapses in rat hippocampus (CA1) at postnatal day 15 and adult ages: implications for the maturation of synaptic physiology and long-term potentiation. *J. Neurosci.* 12, 2685–2705.
- Hayashi, Y., Majewska, A.K., 2005. Dendritic spine geometry: functional implication and regulation. *Neuron* 46, 529–532. doi: [10.1016/j.neuron.2005.05.006](#).
- Heffernan, A.L., Chidgey, C., Peng, P., Masters, C.L., Roberts, B.R., 2016. The neurobiology and age-related prevalence of the ε4 allele of apolipoprotein E in Alzheimer's disease cohorts. *J. Mol. Neurosci.* 60, 316–324. doi: [10.1007/s12031-016-0804-x](#).
- Henderson, V.W., 2006. Estrogen-containing hormone therapy and Alzheimer's disease risk: understanding discrepant inferences from observational and experimental research. *Neuroscience* 138, 1031–1039. doi: [10.1016/j.neuroscience.2005.06.017](#).
- Inagaki, T., Frankfurt, M., Luine, V., 2012. Estrogen-induced memory enhancements are blocked by acute bisphenol A in adult female rats: role of dendritic spines. *Endocrinology* 153, 3357–3367. doi: [10.1210/en.2012-1121](#).
- Ishunina, T.A., Swaab, D.F., 2003. Increased neuronal metabolic activity and estrogen receptors in the vertical limb of the diagonal band of Broca in Alzheimer's disease: relation to sex and aging. *Exp. Neurol.* 183, 159–172. doi: [10.1016/s0014-4886\(03\)00138-9](#).
- Jacobs, E.G., Kroenke, C., Lin, J., Epel, E.S., Kenna, H.A., Blackburn, E.H., Rasgon, N.L., 2013. Accelerated cell aging in female APOE-ε4 carriers: Implications for hormone therapy use. *PLoS One* 8, e54713. doi: [10.1371/journal.pone.0054713](#).
- Jacobs, E.G., Weiss, B.K., Makris, N., Whitfield-Gabrieli, S., Buka, S.L., Klibanski, A., Goldstein, J.M., 2016. Impact of sex and menopausal status on episodic memory circuitry in early midlife. *J. Neurosci.* 36, 10163–10173. doi: [10.1523/JNEUROSCI.0951-16.2016](#).
- Ji, Y., Gong, Y., Gan, W., Beach, T., Holtzman, D.M., Wisniewski, T., 2003. Apolipoprotein E isoform-specific regulation of dendritic spine morphology in apolipoprotein E transgenic mice and Alzheimer's disease patients. *Neuroscience* 122, 305–315. doi: [10.1016/j.neuroscience.2003.08.007](#).
- Joffe, H., Hall, J.E., Gruber, S., Sarmiento, I.A., Cohen, L.S., Yurgelun-Todd, D., Martin, K.A., 2006. Estrogen therapy selectively enhances prefrontal cognitive processes: a randomized, double-blind, placebo-controlled study with functional magnetic resonance imaging in perimenopausal and recently postmenopausal women. *Menopause* 13, 411–422. doi: [10.1097/01.gme.0000189618.48774.7b](#).
- Kang, J.H., Grodstein, F., 2012. Postmenopausal hormone therapy, timing of initiation, APOE and cognitive decline. *Neurobiol. Aging* 33, 1129–1137. doi: [10.1016/j.neurobiolaging.2010.10.007](#).
- Kawas, C., Resnick, S., Morrison, A., Brookmeyer, R., Corrada, M., Zonderman, A., Balcar, C., Lingle, D.D., Metter, E., 1997. A prospective study of estrogen replacement therapy and the risk of developing Alzheimer's disease: the Baltimore Longitudinal Study of Aging. *Neurology* 48, 1517–1521. doi: [10.1212/WNL.48.6.1517](#).
- Kim, J., Szinte, J.S., Boulware, M.I., Frick, K.M., 2016. 17β-estradiol and agonism of G-protein coupled estrogen receptor (GPER) enhance hippocampal memory consolidation via different cell-signaling mechanisms. *J. Neurosci.* 36, 3309–3321. doi: [10.1523/JNEUROSCI.0257-15.2016](#).
- Kim, J., Schalk, J.C., Koss, W.A., Gremminger, R.L., Taxier, L.R., Gross, K.S., Frick, K.M., 2019. Dorsal hippocampal actin polymerization is necessary for activation of G-protein-coupled estrogen receptor (GPER) to increase CA1 dendritic spine density and enhance memory consolidation. *J. Neurosci.* 39, 9598–9610. doi: [10.1523/JNEUROSCI.2687-18.2019](#).
- Koss, W.A., Haertel, J.M., Philippi, S.M., Frick, K.M., 2018. Sex differences in the rapid cell signaling mechanisms underlying the memory-enhancing effects of 17β-estradiol. *eNeuro* 5 doi: [10.1523/ESEURO.0267-18.2018](#).
- Kumar, A., Bean, L.A., Rani, A., Jackson, T., Foster, T.C., 2015. Contribution of estrogen receptor subtypes, ERα, ERβ, and GPER1 in rapid estradiol-mediated enhancement of hippocampal synaptic transmission in mice. *Hippocampus* 25, 1556–1566. doi: [10.1002/hipo.22475](#).
- Kunzler, J., Youmans, K.L., Yu, C., LaDu, M.J., Tai, L.M., 2014. APOE modulates the effect of estrogen therapy on Aβ accumulation EFAD-Tg mice. *Neurosci. Lett.* 560, 131–136. doi: [10.1016/j.neulet.2013.12.032](#).

- Lewis, M.C., Kerr, K.M., Orr, P.T., Frick, K.M., 2008. Estradiol-induced enhancement of object memory consolidation involves NMDA receptors and protein kinase A in the dorsal hippocampus of female C57BL/6 mice. *Behav. Neurosci.* 122, 716–721. doi:[10.1037/0735-7044.122.3.716](https://doi.org/10.1037/0735-7044.122.3.716).
- Li, X.-G., Somogyi, P., Ylinen, A., Buzsáki, G., 1994. The hippocampal CA3 network: an *in vivo* intracellular labeling study. *J. Comp. Neurol.* 339, 181–208. doi:[10.1002/cne.903390204](https://doi.org/10.1002/cne.903390204).
- Liu, D., Pan, X., Zhang, J., Shen, H., Collins, N.C., Cole, A.M., Koster, K.P., Ben Aissa, M., Dai, X., Zhou, M., Tai, L.M., Zhu, Y., LaDu, M.J., Chen, X., 2015. APOE4 enhances age-dependent decline in cognitive function by down-regulating an NMDA receptor pathway in EFAD-Tg mice. *Mol. Neurodegener.* 10, 7. doi:[10.1186/s13024-015-0002-2](https://doi.org/10.1186/s13024-015-0002-2).
- Mahoney, E.R., Dumitrescu, L., Maki, P.M., Rapp, S.R., Keene, C.D., Corrada, M.M.M., Hayden, K.M., Baker, L.D., Espeland, M.A., Resnick, S.M., Hohman, T.J., 2020. Menopausal hormone therapy has beneficial effects on cognitive trajectories among homozygous carriers of the APOE-ε4 allele. *Alzheimers Dement.* 16, e041482. doi:[10.1002/azj.041482](https://doi.org/10.1002/azj.041482).
- Manly, J.J., Merchant, C.A., Jacobs, D.M., Small, S.A., Bell, K., Ferin, M., Mayeux, R., 2000. Endogenous estrogen levels and Alzheimer's disease among postmenopausal women. *Neurology* 54, 833–837. doi:[10.1212/wnl.54.4.833](https://doi.org/10.1212/wnl.54.4.833).
- Mijalkov, M., Volpe, G., Fernaud-Espinosa, I., Defelipe, J., Pereira, J.B., Merino-Serrais, P., 2021. Dendritic spines are lost in clusters in Alzheimer's disease. *Sci. Rep.* 11, 12350. doi:[10.1038/s41598-021-91726-x](https://doi.org/10.1038/s41598-021-91726-x).
- Murakami, G., Tsurugizawa, T., Hatanaka, Y., Komatsuzaki, Y., Tanabe, N., Mukai, H., Hojo, Y., Komimami, S., Yamazaki, T., Kimoto, T., Kawato, S., 2006. Comparison between basal and apical dendritic spines in estrogen-induced rapid spinogenesis of CA1 principal neurons in the adult hippocampus. *Biochem. Biophys. Res. Commun.* 351, 553–558. doi:[10.1016/j.bbrc.2006.10.066](https://doi.org/10.1016/j.bbrc.2006.10.066).
- Nathan, B.P., Barsukova, A.G., Shen, F., McAsey, M., Struble, R.G., 2004. Estrogen facilitates neurite extension via Apolipoprotein E in cultured adult mouse cortical neurons. *Endocrinology* 145, 3065–3073. doi:[10.1210/en.2003-1707](https://doi.org/10.1210/en.2003-1707).
- Packard, M.G., Teather, L.A., 1997. Intra-hippocampal estradiol infusion enhances memory in ovariectomized rats. *NeuroReport* 8, 3009–3013.
- Paganini-Hill, A., Henderson, V.W., 1996. Estrogen replacement therapy and risk of Alzheimer disease. *Arch. Intern. Med.* 156, 2213–2217.
- Phan, A., Suschkov, S., Molinaro, L., Reynolds, K., Lymer, J.M., Bailey, C.D.C., Kow, L.-M., MacLusky, N.J., Pfaff, D.W., Choleris, E., 2015. Rapid increases in immature synapses parallel estrogen-induced hippocampal learning enhancements. *Proc. Natl. Acad. Sci.* 112, 16018–16023. doi:[10.1073/pnas.1522150112](https://doi.org/10.1073/pnas.1522150112).
- Rapp, S.R., Espeland, M.A., Shumaker, S.A., Henderson, V.W., Brunner, R.L., Manson, J.E., Gass, M.L.S., Stefanick, M.L., Lane, D.S., Hays, J., Johnson, K.C., Coker, L.H., Dailey, M., Bowen, D., Investigators, WHIMS, 2003. Effect of estrogen plus progestin on global cognitive function in postmenopausal women: the Women's Health Initiative Memory Study: a randomized controlled trial. *JAMA* 289, 2663–2672. doi:[10.1001/jama.289.20.2663](https://doi.org/10.1001/jama.289.20.2663).
- Roses, A.D., 1996. Apolipoprotein E alleles as risk factors in Alzheimer's disease. *Annu. Rev. Med.* 47, 387–400. doi:[10.1146/annurev.med.47.1.387](https://doi.org/10.1146/annurev.med.47.1.387).
- Selkoe, D.J., 2002. Alzheimer's disease is a synaptic failure. *Science* 298, 789–791. doi:[10.1126/science.1074069](https://doi.org/10.1126/science.1074069).
- Shumaker, S.A., Legault, C., Rapp, S.R., Thal, L., Wallace, R.B., Ockene, J.K., Hendrix, S.L., Jones, B.N., Assaf, A.R., Jackson, R.D., Kotchen, J.M., Wassertheil-Smoller, S., Wactawski-Wende, J., WHIMS Investigators, 2003. Estrogen plus progestin and the incidence of dementia and mild cognitive impairment in postmenopausal women: the Women's Health Initiative Memory Study: a randomized controlled trial. *JAMA* 289, 2651–2662. doi:[10.1001/jama.289.20.2651](https://doi.org/10.1001/jama.289.20.2651).
- Smejkalova, T., Woolley, C.S., 2010. Estradiol acutely potentiates hippocampal excitatory synaptic transmission through a presynaptic mechanism. *J. Neurosci.* 30, 16137–16148. doi:[10.1523/JNEUROSCI.4161-10.2010](https://doi.org/10.1523/JNEUROSCI.4161-10.2010).
- Smith, C.C., Vedder, L.C., McMahon, L.L., 2009. Estradiol and the relationship between dendritic spines, NR2B containing NMDA receptors, and the magnitude of long-term potentiation at hippocampal CA3-CA1 synapses. *Psychoneuroendocrinology* 34 (Suppl 1), S130–S142. doi:[10.1016/j.psyneuen.2009.06.003](https://doi.org/10.1016/j.psyneuen.2009.06.003).
- Spires-Jones, T.L., Meyer-Luehmann, M., Osetek, J.D., Jones, P.B., Stern, E.A., Bacskai, B.J., Hyman, B.T., 2007. Impaired spine stability underlies plaque-related spine loss in an Alzheimer's disease mouse model. *Am. J. Pathol.* 171, 1304–1311. doi:[10.2353/ajpath.2007.070055](https://doi.org/10.2353/ajpath.2007.070055).
- Spruston, N., 2008. Pyramidal neurons: dendritic structure and synaptic integration. *Nat. Rev. Neurosci.* 9, 206–221. doi:[10.1038/nrn2286](https://doi.org/10.1038/nrn2286).
- Tai, L.M., Balu, D., Avila-Munoz, E., Abdullah, L., Thomas, R., Collins, N., Valencia-Olvera, A.C., LaDu, M.J., 2017. EFAD transgenic mice as a human APOE relevant preclinical model of Alzheimer's disease. *J. Lipid Res.* 58, 1733–1755. doi:[10.1194/jlr.R076315](https://doi.org/10.1194/jlr.R076315).
- Takahashi, R.H., Capetillo-Zarate, E., Lin, M.T., Milner, T.A., Gouras, G.K., 2010. Co-occurrence of Alzheimer's disease β-amyloid and tau pathologies at synapses. *Neurobiol. Aging* 31, 1145–1152. doi:[10.1016/j.neurobiolaging.2008.07.021](https://doi.org/10.1016/j.neurobiolaging.2008.07.021).
- Tang, M.-X., Jacobs, D., Stern, Y., Marder, K., Schofield, P., Gurland, B., Andrews, H., Mayeux, R., 1996. Effect of oestrogen during menopause on risk and age at onset of Alzheimer's disease. *The Lancet* 348, 429–432. doi:[10.1016/S0140-6736\(96\)03356-9](https://doi.org/10.1016/S0140-6736(96)03356-9).
- Taxier, L.R., Philippi, S.M., Fortress, A.M., Frick, K.M., 2019. Dickkopf-1 blocks 17β-estradiol-enhanced object memory consolidation in ovariectomized female mice. *Horm. Behav.* 114, 104545. doi:[10.1016/j.yhbeh.2019.06.009](https://doi.org/10.1016/j.yhbeh.2019.06.009).
- Taxier, L.R., Philippi, S.M., York, J.M., LaDu, M.J., Frick, K.M., 2022a. The detrimental effects of APOE4 on risk for Alzheimer's disease may result from altered dendritic spine density, synaptic proteins, and estrogen receptor alpha. *Neurobiol. Aging* 112, 74–86. doi:[10.1016/j.neurobiolaging.2021.12.006](https://doi.org/10.1016/j.neurobiolaging.2021.12.006).
- Taxier, L.R., Philippi, S.M., York, J.M., LaDu, M.J., Frick, K.M., 2022b. APOE4 genotype or ovarian hormone loss influence open field exploration in an EFAD mouse model of Alzheimer's disease. *Horm. Behav.* 140, 105124. doi:[10.1016/j.yhbeh.2022.105124](https://doi.org/10.1016/j.yhbeh.2022.105124).
- Tsolaki, M., Grammaticos, P., Karanasou, C., Balaris, V., Kapoukranidou, D., Kalpidis, I., Petsanis, K., Dedousi, E., 2005. Serum estradiol, progesterone, testosterone, FSH and LH levels in postmenopausal women with Alzheimer's dementia. *Hell. J. Nucl. Med.* 8, 39–42.
- Tuscher, J.J., Luine, V., Frankfurt, M., Frick, K.M., 2016a. Estradiol-mediated spine changes in the dorsal hippocampus and medial prefrontal cortex of ovariectomized female mice depend on ERK and mTOR activation in the dorsal hippocampus. *J. Neurosci.* 36, 1483–1489. doi:[10.1523/JNEUROSCI.3135-15.2016](https://doi.org/10.1523/JNEUROSCI.3135-15.2016).
- Tuscher, J.J., Szinte, J.S., Starrett, J.R., Krentzel, A.A., Fortress, A.M., Remage-Healey, L., Frick, K.M., 2016b. Inhibition of local estrogen synthesis in the hippocampus impairs hippocampal memory consolidation in ovariectomized female mice. *Horm. Behav.* 83, 60–67. doi:[10.1016/j.yhbeh.2016.05.001](https://doi.org/10.1016/j.yhbeh.2016.05.001).
- Tuscher, J.J., Taxier, L.R., Fortress, A.M., Frick, K.M., 2018. Chemogenetic inactivation of the dorsal hippocampus and medial prefrontal cortex, individually and concurrently, impairs object recognition and spatial memory consolidation in female mice. *Neurobiol. Learn. Mem.* 156, 103–116. doi:[10.1016/j.nlm.2018.11.002](https://doi.org/10.1016/j.nlm.2018.11.002).
- Wang, J.M., Irwin, R.W., Brinton, R.D., 2006. Activation of estrogen receptor alpha increases and estrogen receptor beta decreases apolipoprotein E expression in hippocampus *in vitro* and *in vivo*. *Proc. Natl. Acad. Sci. U. S. A.* 103, 16983–16988. doi:[10.1073/pnas.0608128103](https://doi.org/10.1073/pnas.0608128103).
- Whitlock, J.R., Heynen, A.J., Shuler, M.G., Bear, M.F., 2006. Learning induces long-term potentiation in the hippocampus. *Science* 313, 1093–1097. doi:[10.1126/science.1128134](https://doi.org/10.1126/science.1128134).
- Yaffe, K., Haan, M., Byers, A., Tangen, C., Kuller, L., 2000. Estrogen use, APOE, and cognitive decline: evidence of gene-environment interaction. *Neurology* 54, 1949–1954. doi:[10.1212/wnl.54.10.1949](https://doi.org/10.1212/wnl.54.10.1949).
- Youmans, K.L., Tai, L.M., Nwabuisi-Heath, E., Jungbauer, L., Kanekiyo, T., Gan, M., Kim, J., Eimer, W.A., Estus, S., Rebeck, G.W., Weeber, E.J., Bu, G., Yu, C., LaDu, M.J., 2012. APOE4-specific changes in Aβ accumulation in a new transgenic mouse model of Alzheimer disease. *J. Biol. Chem.* 287, 41774–41786. doi:[10.1074/jbc.M112.407957](https://doi.org/10.1074/jbc.M112.407957).
- Yue, X., Lu, M., Lancaster, T., Cao, P., Honda, S.-I., Staufenbiel, M., Harada, N., Zhong, Z., Shen, Y., Li, R., 2005. Brain estrogen deficiency accelerates Aβ plaque formation in an Alzheimer's disease animal model. *Proc. Natl. Acad. Sci.* 102, 19198–19203. doi:[10.1073/pnas.0505203102](https://doi.org/10.1073/pnas.0505203102).