

# INVARIANT DIFFERENTIAL OPERATORS AND FCR FACTORS OF ENVELOPING ALGEBRAS

IAN M. MUSSON AND JEB F. WILLENBRING

ABSTRACT. If  $\mathfrak{g}$  is a semisimple Lie algebra, we describe the prime factors of  $\mathcal{U}(\mathfrak{g})$  that have enough finite dimensional modules. The proof depends on some combinatorial facts about the Weyl group which may be of independent interest. We also determine, which finite dimensional  $\mathcal{U}(\mathfrak{g})$ -modules are modules over a given prime factor. As an application we study finite dimensional modules over some rings of invariant differential operators arising from Howe duality.

## 1. INTRODUCTION

Let  $F$  denote an algebraically closed field of characteristic zero. Given an  $F$ -algebra  $A$ , we say that  $A$  *has enough finite dimensional representations* if the intersection of the annihilators of the finite dimensional  $A$ -modules is zero. If in addition all finite dimensional representations are completely reducible  $A$  is an *FCR* algebra (see [KS94]).

Let  $\mathfrak{g}$  be a semisimple Lie algebra with enveloping algebra  $\mathcal{U}(\mathfrak{g})$ . In this paper, we describe the prime factors of  $\mathcal{U}(\mathfrak{g})$  that are FCR algebras, and also determine, in terms of highest weights, which finite dimensional  $\mathcal{U}(\mathfrak{g})$ -modules are actually modules over a given prime factor. If  $A$  is a prime factor of  $\mathcal{U}(\mathfrak{g})$ , then any finite dimensional  $A$ -module is a  $\mathfrak{g}$ -module, and hence completely reducible. So the issue is whether or not  $A$  has enough finite dimensional modules. However we use the appellation “FCR” for brevity.

We begin with some basic notation used to state the main theorem. Suppose that  $\mathfrak{g}$  is a reductive Lie algebra. Let  $\mathfrak{g}_{ss} = [\mathfrak{g}, \mathfrak{g}]$ , be the semisimple part of  $\mathfrak{g}$ . Fix a Cartan subalgebra  $\mathfrak{h}$  of  $\mathfrak{g}$  and set  $\mathfrak{h}_{ss} = \mathfrak{h} \cap \mathfrak{g}_{ss}$ . Then,  $\mathfrak{h} = \mathfrak{h}_{ss} \oplus \mathfrak{z}$  where  $\mathfrak{z}$  is the center of  $\mathfrak{g}$ .

Now we recall some results of Soergel on prime ideals in the enveloping algebra  $\mathcal{U} = \mathcal{U}(\mathfrak{g})$ , for  $\mathfrak{g}$  reductive. To any prime ideal  $\Omega$  of  $\mathcal{S} = \mathcal{S}(\mathfrak{h})$ , the symmetric algebra on  $\mathfrak{h}$ , we can associate a prime ideal  $I_\Omega$  of  $\mathcal{U}$ , and a “tautological highest weight”  $\lambda := \lambda_\Omega$  (see [Soe90]). We recall the details in Section 3. Any prime ideal of  $\mathcal{U}$  has the form  $I_\Omega$  for a suitable prime ideal  $\Omega$  in  $\mathcal{S}$ . Set  $\mathcal{U}_\Omega := \mathcal{U}/I_\Omega$ .

Let  $R$  denote the root system corresponding to the pair  $(\mathfrak{g}, \mathfrak{h})$  with Weyl group  $W$ . Choose a set of positive roots,  $R^+$ , such that  $R = R^+ \cup -R^+$ , and let  $B = \{\alpha_1, \alpha_2, \dots, \alpha_n\}$  denote the simple roots in  $R^+$ .

Set  $\rho := \frac{1}{2} \sum_{\alpha \in R^+} \alpha$ , and define the “dot action” of  $W$  on  $\mathfrak{h}^*$  as:

$$w.\xi := w(\xi + \rho) - \rho \quad \text{for } \xi \in \mathfrak{h}^* \text{ and } w \in W.$$

---

*Date:* September 19, 2006.

The first author was supported in part by NSF Grant DMS-0099923.

The second author was supported in part by NSA Grant H98230-05-1-0078.

The dot action of  $W$  on  $\mathfrak{h}^*$  induces an action on  $\mathcal{S}$ . Let  $\kappa(\cdot, \cdot)$  denote the Killing form on  $\mathfrak{g}_{ss}$ . For  $\mu \in \mathfrak{h}^*$ , let  $h_\mu$  be the element of  $\mathfrak{h}_{ss}$  determined by  $\kappa(h_\mu, h) = \mu(h)$  for all  $h \in \mathfrak{h}_{ss}$ . The isomorphism  $\mathfrak{h}_{ss}^* \simeq \mathfrak{h}_{ss}$  sending  $\mu$  to  $h_\mu$  induces a bilinear form  $(\cdot, \cdot)$  on  $\mathfrak{h}_{ss}^*$ . Recall that this form is positive definite on the real span of the roots. For  $\alpha \in R$  let  $H_\alpha = \frac{2h_\alpha}{(\alpha, \alpha)}$  denote the coroot to  $\alpha$ , and set  $\alpha^\vee = \frac{2\alpha}{(\alpha, \alpha)}$ . Let  $\bar{\omega}_i \in \mathfrak{h}_{ss}^*$  be the fundamental weights, so that  $(\bar{\omega}_i, \alpha_j^\vee) = \delta_{i,j}$ . We write  $H_i$  (resp.  $h_i$ ) in place of  $H_{\alpha_i}$  (resp.  $h_{\alpha_i}$ ). Thus  $\mu(H_i) = (\mu, \alpha_i^\vee)$ .

Each  $\lambda \in \mathfrak{h}^*$  determines a subroot system,  $R_\lambda = \{\alpha \in R \mid (\lambda, \alpha^\vee) \in \mathbb{Z}\}$ . Let  $W_\lambda$  denote the Weyl group of  $R_\lambda$ . Let  $B_\lambda$  be the unique basis of simple roots in  $R_\lambda^+ := R^+ \cap R_\lambda$ , and set  $W^\lambda := \{w \in W \mid w(B_\lambda) \subseteq R^+\}$ . The elements of  $W^\lambda$  are left coset representatives for  $W_\lambda$  in  $W$ . Thus,  $W = W^\lambda W_\lambda$ .

For  $\mu \in \mathfrak{h}^*$ , let  $L(\mu)$  be the simple module with highest weight  $\mu$ . Let  $P^+ = \{\mu \in \mathfrak{h}^* \mid \dim L(\mu) < \infty\}$ . We say that  $\mu$  is *dominant* if  $\mu \in P^+$ . It is a result of Harish-Chandra [HC] that  $\mathcal{U}$  has enough finite dimensional representations, that is  $\bigcap_{\lambda \in P^+} \text{ann} L(\lambda) = 0$ . This fact, together with the fact that  $P^+$  is dense in  $\mathfrak{h}^*$  can be used to show the injectivity of the Harish-Chandra homomorphism (see [Jo98] Section 7). Thus it is not surprising that we need a notion of dominance for prime ideals. We say that a prime ideal  $\Omega$  is *dominant* if  $\mathcal{V}(\Omega) \cap P^+$  is Zariski dense in  $\mathcal{V}(\Omega)$ , where  $\mathcal{V}(\Omega)$  denotes the zero set of  $\Omega$  in  $\mathfrak{h}^*$ .

What is surprising, perhaps, is that it is not necessary for  $\Omega$  to be dominant for  $\mathcal{U}_\Omega$  to be FCR.

**Main Theorem.** *Assume  $\mathfrak{g}$  is semisimple and  $\mathcal{U} = \mathcal{U}(\mathfrak{g})$ .*

- (1) *If  $\mathcal{U}_\Omega$  is FCR then for some  $w \in W$ ,  $w.\Omega$  is dominant.*
- (2) *Let  $\Omega$  be a dominant ideal in  $\mathcal{S}$ , and  $w \in W$ .*

*The following conditions are equivalent:*

- (a)  $\mathcal{U}_{w.\Omega}$  is FCR
- (b)  $I_{w.\Omega} = I_\Omega$
- (c)  $w \in W^\lambda$  where  $\lambda = \lambda_\Omega$ .

This paper is organized as follows. In Section 2 we prove some results about root systems and the Weyl group which may be of independent interest. Then in Section 3 we collect some results about prime and primitive ideals in enveloping algebras. The main theorem is proved in Section 4. In Sections 5, we apply the main theorem to some examples of rings of invariant differential operators related to Howe duality.

To end this introduction, we mention that further results on prime ideals in  $\mathcal{U}$  are obtained in [BJ01], [P05], and [P].

## 2. WEYL GROUP COMBINATORICS

We introduce some notation. Let  $B_1$  be a subset of  $B$ ,  $R_1$  be the subroot system of  $R$  with simple roots  $B_1$  and let  $R_1^+$  be the corresponding set of positive roots. Let  $W_1$  be the Weyl group of  $R_1$  and set

$$\rho' = \frac{1}{2} \sum_{\alpha \in R_1^+} \alpha.$$

For  $w \in W$  define:

$$Q(w) = \{\alpha \in R^+ \mid w\alpha \in -R^+\},$$

and set  $\ell(w) = |Q(w)|$ . We have ([GW98] Lemma 7.3.6),

$$\rho - w^{-1}\rho = \sum_{\alpha \in Q(w)} \alpha. \quad (1)$$

**Lemma 2.1.** *Suppose  $w \in W$  is such that  $w(R_1) = R_1$  and set*

$$T(w) = \{\alpha \in R_1^+ | w\alpha \in -R_1^+\}.$$

*Then,*

$$\rho' - w^{-1}\rho' = \sum_{\alpha \in T(w)} \alpha.$$

*Proof.* Note that  $R_1^+$  is a disjoint union  $R_1^+ = w(R_1^+ \setminus T(w)) \cup (-wT(w))$ . The rest of the proof is similar to the proof of (1).  $\square$

**Proposition 2.2.** *Suppose that  $w \in W$  and  $w(R_1) = R_1$ . Then*

$$(\rho', \sum_{\alpha \in Q(w)} \alpha) \geq 0,$$

*with equality if and only if  $w(B_1) = B_1$ .*

We introduce some notation needed for the proof of this result. If  $Q$  is a finite subset of  $\mathfrak{h}^*$  we set  $\langle Q \rangle = \sum_{\alpha \in Q} \alpha$ . Let

$$W_0 = \{w \in W | w(R_1) = R_1\},$$

and

$$T = \{t \in W | t(B_1) = B_1\}.$$

**Lemma 2.3.** *For  $w \in W_0$  the following are equivalent,*

- (1)  $w(B_1) = B_1$ ,
- (2)  $w(B_1) \subseteq R_1^+$ ,
- (3)  $\ell(ws_\alpha) > \ell(w)$  for all  $\alpha \in B_1$ .

*Proof.* Clearly (1) implies (2). For the reverse implication, suppose  $\beta \in R_1^+$  and write  $\beta = \sum_{\alpha \in B_1} c_\alpha \alpha$ . Define  $ht(\beta) = \sum_{\alpha} c_\alpha$ . From (2) it follows that  $ht(w(\beta)) \geq ht(\beta)$  and  $w(R_1^+) = R_1^+$ . This implies that  $w^{-1}\gamma \in B_1$  for  $\gamma \in B_1$ . Hence (1) holds. The equivalence of (2) and (3) follows from [Hu90] Lemma 1.6 a), b).  $\square$

**Corollary 2.4.** *Set  $T = \{w \in W | w(B_1) = B_1\}$ . Then,*

- (1)  $W_0 = W_1 T$  is the semidirect product of the normal subgroup  $W_1$  by  $T$ .
- (2) For  $w \in W_1$  and  $t \in T$  we have  $\ell(wt) = \ell(w) + \ell(t)$ .

*Proof.* (1) Since  $W_1$  is generated by reflections,  $s_\alpha$  with  $\alpha \in B_1$  and  $ts_\alpha t^{-1} = s_{t(\alpha)}$  for  $t \in T$ , it follows that  $T$  normalizes  $W_1$ . From the implication (1)  $\implies$  (3) in Lemma 2.3 and [Hu90] Proposition 1.10 (c) we see that  $T$  is a transversal to  $W_1$  in  $W_0$ . It follows easily that  $W_0 = W_1 T$  is the semidirect product of  $W_1$  by  $T$ .

(2) This holds by [Hu90] Proposition 1.10 (c).  $\square$

Recall that if  $w \in W$  and  $\alpha \in B$  such that  $\ell(ws_\alpha) = \ell(w) + 1$  we have  $Q(ws_\alpha) = s_\alpha Q(w) \cup \{\alpha\}$  ([GW98] Corollary 7.3.4). If  $w_1, w_2 \in W$  and  $\ell(w_1 w_2) = \ell(w_1) + \ell(w_2)$ , it follows by induction on  $\ell(w_2)$  that  $Q(w_1 w_2) = w_2^{-1} Q(w_1) \cup Q(w_2)$ , a disjoint

union. Now for  $t \in T$  and  $w \in W_1$  we have  $\ell(wt) = \ell(w) + \ell(t)$  by Corollary 2.4. Since  $t\rho' = \rho'$ , it follows that

$$(\rho', \langle Q(wt) \rangle) = (\rho', \langle Q(w) \rangle) + (\rho', \langle Q(t) \rangle).$$

Hence Proposition 2.2 follows from the following result.

Let  $v$  be the longest element of  $W_1$ . Then,  $v(R_1^+) = -R_1^+$  and since  $\ell(v) = |R_1^+|$ , we have  $v(R^+ \setminus R_1^+) = R^+ \setminus R_1^+$ . Note that  $v^2 = 1$ , and if  $t \in T$ , then since  $t^{-1}vt \in W_1$  and  $t^{-1}vt(R_1^+) = -R_1^+$  we have  $tv = vt$ . Suppose  $t \in T$ , and set

$$Q^+ = \{\alpha \in Q(t) \mid (\rho', \alpha) > 0\},$$

$$Q^- = \{\alpha \in Q(t) \mid (\rho', \alpha) < 0\}.$$

**Lemma 2.5.**

- (1) If  $w \in W_1$  then  $Q(w) = T(w)$ . Hence, if  $w \neq 1$ , then  $(\rho', \langle Q(w) \rangle) > 0$ .
- (2) The map  $\kappa : \alpha \mapsto v\alpha$  is a permutation of  $Q(t)$ , which interchanges  $Q^+$  and  $Q^-$ .
- (3) If  $t \in T$ , then  $(\rho', \langle Q(t) \rangle) = 0$ .

*Proof.* (1) Obviously  $T(w) \subseteq Q(w)$ , and it follows from [Hu90] Corollary 1.7 and Proposition 1.10 (b) that  $\ell(w) = |T(w)|$ , giving equality. The second statement follows from ([GW98] Lemma 2.5.12).

(2) First observe that if  $\alpha \in Q(t)$  then  $\alpha \notin R_1^+$ , since  $t(R_1^+) = R_1^+$  hence  $t\alpha = -\beta$  with  $\beta \in R^+ \setminus R_1^+$ . We must show that  $v\alpha$  is in  $R^+$  and  $tv\alpha \in -R^+$ . The first statement follows since  $v(R^+ \setminus R_1^+) = R^+ \setminus R_1^+$ , and the second follows from  $tv\alpha = vt\alpha = -v\beta \in -R^+$ . We have shown that  $\kappa$  is a permutation of  $Q(t)$ . Finally, for  $\alpha \in Q(t)$ ,

$$(\rho', v\alpha) = (v\rho', \alpha) = -(\rho', \alpha).$$

The result follows easily from this fact.

(3) follows immediately from (2).  $\square$

Next we relate the dot action of  $W$  on  $\mathcal{S}$  to the usual action.

**Lemma 2.6.** If  $h \in \mathfrak{h}$  and  $w \in W$  then,

$$w.h = wh - \sum_{\alpha \in Q(w)} h(\alpha).$$

*Proof.* For  $\lambda \in \mathfrak{h}^*$  we have,

$$\begin{aligned} (w.h)(\lambda) &= h(w^{-1}.\lambda) \\ &= h(w^{-1}(\lambda + \rho) - \rho) \\ &= (wh)(\lambda) + h(w^{-1}\rho - \rho). \end{aligned}$$

We now apply equation (1).  $\square$

### 3. FCR FACTORS OF ENVELOPING ALGEBRAS.

We recall Soergel's work on prime ideals in the enveloping algebra  $\mathcal{U} = \mathcal{U}(\mathfrak{g})$  where  $\mathfrak{g}$  is a reductive Lie algebra. Let  $\mathcal{Z}$  denote the center of  $\mathcal{U}$ .

If  $R$  is a ring we let  $\text{Spec}(R)$  denote the prime spectrum of  $R$ . For  $\Omega \in \text{Spec } \mathcal{S}$ , define an ideal  $I_\Omega$  in  $\text{Spec } \mathcal{U}$  as follows: Let  $\mathbb{F} := \text{Quot}(\mathcal{S}/\Omega)$  (the quotient field of the commutative domain defined by  $\Omega$ ). We will let  $\mathfrak{g}_{\mathbb{F}} = \mathbb{F} \otimes_{\mathbb{F}} \mathfrak{g}$  denote the Lie

algebra obtained from  $\mathfrak{g}$  by extension of scalars. We write  $\lambda_\Omega$  for the  $\mathbb{F}$ -linear map from  $\mathfrak{h}_\mathbb{F}$  to  $\mathbb{F}$  whose restriction to  $\mathfrak{h}$  is:

$$\lambda_\Omega : \mathfrak{h} \hookrightarrow \mathcal{S} \twoheadrightarrow \mathcal{S}/\Omega \hookrightarrow \mathbb{F}.$$

Let  $L(\lambda_\Omega)$  be the simple highest weight module for  $\mathfrak{g}_\mathbb{F}$  with highest weight  $\lambda_\Omega$ . Let  $I_\Omega$  be the annihilator of  $L(\lambda_\Omega)$  in  $\mathcal{U}(\mathfrak{g})$  and set  $\phi(\Omega) = I_\Omega$ . The maximal ideals in  $\mathcal{S}$  have the form,  $M_\mu := \{f \in \mathcal{S} | f(\mu) = 0\}$  for  $\mu \in \mathfrak{h}^*$ , and we write  $I_\mu$  instead of  $I_{M_\mu}$ , so that  $I_\mu$  is the annihilator of the simple module  $L(\mu)$  with highest weight  $\mu$ . We write  $\mathfrak{h}_\mathbb{F}^*$  for  $\mathfrak{h}^* \otimes \mathbb{F} = (\mathfrak{h}_\mathbb{F})^*$ .

*Remark 3.1.* When  $\lambda = \lambda_\Omega$  we have:

$$R_\lambda = \{\alpha \in R | H_\alpha - n \in \Omega \text{ for some } n \in \mathbb{Z}\}.$$

For  $P \in \text{Spec } \mathcal{U}$ , let  $\psi(P) = P \cap Z$  ( $\in \text{Spec } \mathcal{Z}$ ). Lastly,  $\theta$  denotes the map induced by the Harish-Chandra isomorphism  $\mathcal{Z} \xrightarrow{\sim} \mathcal{S}^W \subseteq \mathcal{S}$ . Then (by [Soe90], Section 2.1) we have the following commutative diagram:

$$\begin{array}{ccc} \text{Spec } \mathcal{S} & & \\ \downarrow \theta & \searrow \phi & \\ & \text{Spec } \mathcal{U} & \\ & \swarrow \psi & \\ \text{Spec } \mathcal{Z} & & \end{array}$$

By [Soe90] Section 2.2, these maps are continuous in the Jacobson topology on prime ideals. In particular these maps preserve inclusions.

Note that the sets  $R_\lambda$ ,  $W_\lambda$ ,  $B_\lambda$  depend only on the coset  $\Lambda := \lambda + P(R) \in \mathfrak{h}_\mathbb{F}^*/P(R)$ . For a coset  $\Lambda$ , let  $B_\Lambda = B_\lambda$  for any  $\lambda \in \Lambda$ .

Set

$$\Lambda^+ = \{\lambda \in \Lambda | (\lambda + \rho, \alpha^\vee) \geq 0, \forall \alpha \in B_\Lambda\}, \text{ and}$$

$$\Lambda^{++} = \{\lambda \in \Lambda | (\lambda + \rho, \alpha^\vee) > 0, \forall \alpha \in B_\Lambda\}.$$

**Lemma 3.2** ([Jan83] Satz 5.16). *If  $w \in W^\lambda$  then for all  $\mu \in \lambda + P(R)$  we have  $I_\mu = I_{w.\mu}$*

We recall the definition of the  $\tau$ -invariant for primitive ideals. Set

$$\tau_\Lambda(w) = \{\alpha \in B_\Lambda | w.\alpha \in -R^+\}.$$

Next define  $X_\lambda = \{I_{w.\lambda} | w \in W\}$ , and write  $2^{B_\Lambda}$  for the poset of subsets of  $B_\Lambda$  ordered by inclusion. We have:

**Theorem 3.3** ([Jan83] Satz 5.7). *Let  $\Lambda \in \mathfrak{h}^*/P(R)$  and  $\lambda \in \Lambda^{++}$ . Then there is a well defined order reversing map:*

$$\tau_\Lambda : X_\lambda \rightarrow 2^{B_\Lambda}$$

*such that  $\tau_\Lambda(I_{w.\lambda}) = \tau_\Lambda(w)$ .*

**Lemma 3.4.** *Let  $D$  be a dense subset of an affine algebraic set  $X$ . Let:*

$$X = X_1 \cup X_2 \cup X_3 \cup \cdots \cup X_r$$

*be the decomposition of  $X$  into irreducible components. Then  $D \cap X_i$  is dense in  $X_i$  for each  $i$ .*

*Proof.* Consider  $i$  such that  $1 \leq i \leq r$ . There exists a function  $p_i \in \mathcal{O}(X)$  which vanishes on  $X_j$  for all  $j \neq i$  and  $p_i$  is not identically zero on  $X_i$ . Let  $f \in \mathcal{O}(X_i)$  be a function that vanishes on  $D \cap X_i$ . We will show that  $f$  is identically zero on  $X_i$ . We may extend  $f$  to a function  $g \in \mathcal{O}(X)$ . The function  $gp_i$  vanishes on  $D$  and so is identically zero on  $X$ . On  $X_i$ ,  $gp_i = fp_i$ . This means that  $f$  vanishes on the set  $D' = \{x \in X_i | p_i(x) \neq 0\}$ , which is dense in  $X_i$ . Therefore,  $f$  vanishes on  $X_i$ .  $\square$

#### 4. PROOF OF THE MAIN RESULT.

Until further notice  $\mathfrak{g}$  will be a reductive Lie algebra over  $F$ . In general, it is unknown when a prime ideal  $I_\Omega$  is contained in a given primitive ideal  $I_\lambda$ . However for  $\lambda \in P^+$  the answer is easy.

**Theorem 4.1.** *For  $\mu \in P^+$ ,  $I_\Omega \subseteq I_\mu$  if and only if  $W.\mu \cap \mathcal{V}(\Omega) \neq \emptyset$ .*

*Proof.* By Harish-Chandra's theorem we have  $\mathcal{Z} \cong \mathcal{S}^W$ . The maximal ideals in  $\mathcal{S}^W$  are in one-to-one correspondence with points in the orbit space,  $\mathfrak{h}^*/W$ . The embedding  $\mathcal{S}^W \hookrightarrow \mathcal{S}$  induces the canonical map,  $\mathfrak{h}^* \rightarrow \mathfrak{h}^*/W$ , which we write as  $\pi(\mu) = [\mu]$ . This map agrees with the restriction of  $\theta$  to the set of maximal ideals in  $\mathcal{S}$ .

( $\Rightarrow$ ): Suppose  $I_\Omega \subseteq I_\mu$ . Then,  $\theta(\Omega) = \psi(I_\Omega) \subseteq \psi(I_\mu) = \theta(M_\mu)$ . Applying  $\mathcal{V}(\cdot)$  reverses inclusions, so we have  $\mathcal{V}(\theta(\Omega)) \supseteq \mathcal{V}(\theta(M_\mu))$ . By definition,  $[\mu] = \mathcal{V}(\theta(M_\mu))$ . That is to say that there exists  $w \in W$  such that  $w.\mu \in \mathcal{V}(\Omega)$ .

( $\Leftarrow$ ): On the other hand if  $w.\mu \in \mathcal{V}(\Omega)$ , then  $I_\Omega \subseteq I_{w.\mu}$ . This implies, by a result of Dixmier [Dix70] (or [Jan83] Korollar 5.21) that  $I_\Omega \subseteq I_\mu$ .  $\square$

**Definition 4.2.** Set  $\Lambda(\Omega) := \{\mu \in P^+ | I_\Omega \subseteq I_\mu\}$ , and for each  $w \in W$ , set  $\Lambda(\Omega, w) := \{\mu \in P^+ | w.\mu \in \mathcal{V}(\Omega)\}$ .

Note that,  $I_\Omega \subseteq \bigcap_{\mu \in \Lambda(\Omega)} I_\mu$  and equality holds if and only if  $U_\Omega$  is FCR. Also, by Theorem 4.1 we have:

$$\Lambda(\Omega) = \bigcup_{w \in W} \Lambda(\Omega, w).$$

**Proposition 4.3.** *If  $\Omega$  is dominant then  $U_\Omega$  has enough finite dimensional modules.*

*Proof.* If  $P^+ \cap \mathcal{V}(\Omega)$  is dense in  $\mathcal{V}(\Omega)$  then  $I_\Omega = \bigcap_{\mu \in P^+ \cap \mathcal{V}(\Omega)} I_\mu$  by [Soe90] Proposition 1. Thus by the remark after Definition 4.2, it suffices to show that  $P^+ \cap \mathcal{V}(\Omega) \subseteq \Lambda(\Omega)$  because this will imply  $\bigcap_{\mu \in \Lambda(\Omega)} I_\mu \subseteq \bigcap_{\mu \in P^+ \cap \mathcal{V}(\Omega)} I_\mu$ . The former inclusion follows since if  $\mu \in P^+ \cap \mathcal{V}(\Omega)$  then  $\Omega \subseteq M_\mu$ , so  $I_\Omega \subseteq I_\mu$ , because  $\phi$  preserves inclusions.  $\square$

**Lemma 4.4.** *If  $\Omega$  is a dominant ideal in  $\mathcal{S}$ , and  $w \in W^\lambda$  where  $\lambda = \lambda_\Omega$  then  $I_{w.\Omega} = I_\Omega$ .*

*Proof.* If  $w \in W^\lambda$ , then  $I_\lambda = I_{w.\lambda}$  by Lemma 3.2. Thus,  $I_\Omega = I_{w.\Omega}$  by [Soe90] Theorem 1 part (ii).  $\square$

For the remainder of this section we assume that  $\mathfrak{g}$  is semisimple.

**Proposition 4.5.** *If  $\Omega$  is dominant and  $\lambda = \lambda_\Omega$ , then  $B_\lambda \subset B$ .*

*Proof.* Let  $B = \{\alpha_1, \dots, \alpha_n\}$ . It suffices to show that if  $\alpha \in R_\lambda \cap R^+$  and  $\alpha = \sum_{i=1}^n c_i \alpha_i$  ( $c_i \in \mathbb{Z}, c_i \geq 0$ ) then  $c_i > 0$  implies  $\alpha_i \in R_\lambda$ . Now  $\alpha \in R_\lambda$  means  $H_\alpha - m' \in \Omega$  for some  $m' \in \mathbb{Z}$ . If  $\mu \in \mathcal{V}(\Omega) \cap P^+$  we have  $\mu_j = \mu(H_j) \in \mathbb{N}$  for  $1 \leq j \leq n$ . Also,  $h_\alpha = \sum c_j h_j$ . We can write  $H_\alpha = \sum \frac{r_j}{s} H_j$  where  $s, r_j \in \mathbb{N}$ ,  $s > 0$  are such that  $\frac{r_j}{s} = c_j \frac{(\alpha_j, \alpha)}{(\alpha, \alpha)}$ . Then,  $m' = \mu(H_\alpha) = \sum \frac{r_j}{s} \mu_j$ . Hence, if  $c_i > 0$ , then  $r_i > 0$  and  $0 \leq \mu_i \leq \frac{m's}{r_i} = m$ .

Suppose  $c_i > 0$ , and for  $0 \leq p \leq m$ , let  $\Pi_{i,p} := \mathcal{V}(H_i - p)$ . The above establishes that  $\mathcal{V}(\Omega) \cap P^+ \subseteq \bigcup_{p=0}^m \Pi_{i,p}$ . Since  $\Omega$  is dominant,  $\mathcal{V}(\Omega) \subseteq \bigcup_{p=0}^m \Pi_{i,p}$ , but  $\mathcal{V}(\Omega)$  is irreducible so  $\mathcal{V}(\Omega) \subseteq \Pi_{i,p}$  for some  $p$ . This means  $H_i - p \in \Omega$ , so  $\alpha_i \in R_\lambda$ .  $\square$

To prove an important special case of the main theorem. Before proceeding however, observe that for any ideal  $\Omega \subseteq \mathcal{S}$  and  $w \in W$  we have:

$$\mathcal{V}(w\Omega) = w\mathcal{V}(\Omega), \quad \text{and} \quad \mathcal{V}(w.\Omega) = w.\mathcal{V}(\Omega).$$

Furthermore, if  $\Omega'$  is an ideal such that  $\mathcal{V}(\Omega') = \mathcal{V}(\Omega) + \rho$ , then for any  $w \in W$ ,  $\mathcal{V}(w\Omega') = w.\mathcal{V}(\Omega) + \rho$ . In addition  $H_\alpha - n \in \Omega$  implies  $H_\alpha - n - (\rho, \alpha^\vee) \in \Omega'$ .

**Theorem 4.6.** *Suppose  $\Omega$  and  $w.\Omega$  are prime ideals of  $\mathcal{S}$  such that  $P^+ \cap \mathcal{V}(\Omega)$  and  $P^+ \cap \mathcal{V}(w.\Omega)$  are nonempty. Then  $w \in W^\lambda$ , where  $\lambda := \lambda_\Omega$  is the tautological highest weight corresponding to  $\Omega$ .*

*Proof.* Suppose that  $\mu \in P^+ \cap \mathcal{V}(\Omega)$  and  $\nu \in P^+ \cap \mathcal{V}(w.\Omega)$ . Assume to the contrary that  $w \in W$  is such that there exists  $\alpha \in B_\lambda$  such that  $w\alpha = -\beta$  for  $\beta \in R^+$ . Since  $\alpha \in B_\lambda$  we have  $H_\alpha - n \in \Omega$  for some  $n \in \mathbb{Z}$ . Because  $\mu \in P^+ \cap \mathcal{V}(\Omega)$ , it follows that  $\mu(H_\alpha - n) = 0$ , and  $\mu(H_\alpha) \in \mathbb{N}$ . This implies that  $n \in \mathbb{N}$ . Similarly  $H_\beta - m \in w.\Omega$  for some  $m \in \mathbb{N}$ .

Let  $\Omega'$  be the ideal in  $\mathcal{S}$  defined by the closed algebraic set  $\mathcal{V}(\Omega) + \rho$ . Then  $H_\alpha - n - (\rho, \alpha^\vee) \in \Omega'$ , and so  $-w(H_\alpha - n - (\rho, \alpha^\vee)) = H_\beta + n + (\rho, \alpha^\vee) \in w\Omega'$ . Since  $\nu \in \mathcal{V}(w.\Omega) = w.\mathcal{V}(\Omega) = \mathcal{V}(w\Omega') - \rho$ , we have,  $\nu + \rho \in \mathcal{V}(w\Omega')$ . Therefore,

$$(\nu + \rho)(H_\beta + n + (\rho, \alpha^\vee)) = 0,$$

since an element of  $w\Omega'$  evaluates as 0 on  $\mathcal{V}(w\Omega')$ . This means that:

$$m + (\rho, \beta^\vee) + n + (\rho, \alpha^\vee) = 0,$$

But this is a contradiction because  $(\rho, \gamma^\vee) > 0$  for all  $\gamma \in R^+$ .  $\square$

In the next two lemmas we assume that  $\Omega$  is dominant and  $\lambda = \lambda_\Omega$ . By Proposition 4.5,  $B_\lambda \subseteq B$ . Thus, we may apply the results of Section 2 with  $B_1$  replaced by  $B_\lambda$  etc. Let  $I$  the subset of  $\{1, 2, \dots, \ell\}$  such that  $B_\lambda = \{\alpha_i | i \in I\}$ .

**Lemma 4.7.** *Assume that  $\Omega$  is dominant and  $\lambda = \lambda_\Omega$ . Then if  $w \in W$ ,  $w(R_\lambda) = R_\lambda$  and  $w(B_\lambda) \not\subseteq R_\lambda^+$  then*

$$w.h_{\rho'} = wh_{\rho'} - c.$$

for some  $c > 0$ .

*Proof.* By Lemma 2.6,

$$w.h_{\rho'} = wh_{\rho'} - h_{\rho'} \left( \sum_{\alpha \in Q(w)} \alpha \right).$$

Since  $\Omega$  is dominant  $B_\lambda \subseteq B$  by Proposition 4.5. Therefore by Proposition 2.2,  $c := h_{\rho'}(\sum_{\alpha \in Q(w)} \alpha) > 0$ .  $\square$

**Lemma 4.8.** *If  $w \in W$  and  $w.\Omega = \Omega$  then  $w(B_\lambda) = B_\lambda$ .*

*Proof.* Suppose that  $w.\Omega = \Omega$ . We proceed in two steps.

Step 1.  $w(R_\lambda) = R_\lambda$ : Indeed, if  $\alpha \in R_\lambda$ , then  $H_\alpha - m \in \Omega$  for some  $m \in \mathbb{Z}$ . If  $w\alpha = \beta$  then by Lemma 2.6,  $\Omega$  contains

$$w.(H_\alpha - m) = H_\beta - \sum_{\gamma \in Q(w)} (\gamma, \alpha^\vee) - m,$$

and this implies that  $\beta \in R_\lambda$ .

Step 2: Since the  $\alpha_i$  ( $i \in I$ ) belong to  $R_\lambda$  there are non-negative rational numbers  $a_i$  ( $i \in I$ ) such that:

$$h_i - a_i \in \Omega$$

similarly,  $h_{\rho'} - a \in \Omega$  for some non-negative  $a \in \mathbb{Q}$ . By Lemma 2.1 we have

$$w\rho' - \rho' = \sum_{\alpha \in T(w)} w\alpha = - \sum_{i \in I} b_i \alpha_i$$

for non-negative integers  $b_i$ . Hence,

$$wh_{\rho'} = h_{\rho'} - \sum_{i \in I} b_i h_i. \quad (2)$$

Assume for a contradiction that  $w(B_\lambda) \not\subseteq B_\lambda$ . Then by Lemma 2.3  $w(B_\lambda) \not\subseteq R_\lambda^+$ , so by Lemma 4.7,

$$w.h_{\rho'} = wh_{\rho'} - c \quad (3)$$

for some  $c > 0$ . Since  $w.\Omega = \Omega$ , it follows from equations (2) and (3) that  $\Omega$  contains,

$$\begin{aligned} w.(h_{\rho'} - a) &= wh_{\rho'} - a - c \\ &= h_{\rho'} - \sum_{i \in I} b_i h_i - a - c. \end{aligned}$$

However,  $h_i - a_i \in \Omega$  for  $i \in I$ , so

$$h_{\rho'} - \sum_{i \in I} b_i a_i - a - c \in \Omega.$$

Since  $h_{\rho'} - a \in \Omega$  we deduce that

$$\sum_{i \in I} b_i a_i + c \in \Omega.$$

This is a contradiction since  $c > 0$  and  $b_i, a_i \geq 0$  for all  $i \in I$ .  $\square$

**Theorem 4.9.** *If  $U_{\Omega'}$  is FCR there exists a dominant ideal  $\Omega$  in  $\mathcal{S}$  such that  $\Omega' = w.\Omega$  for some  $w \in W^\lambda$  (where  $\lambda := \lambda_\Omega$  is the tautological highest weight of  $\Omega$ ).*



*Proof.* Let

$$X := \overline{\Lambda(\Omega')} = X_1 \cup \cdots \cup X_r.$$

define the decomposition of  $X$  into irreducible components. Set  $Y_i := \Lambda(\Omega') \cap X_i$ . Since  $U_{\Omega'}$  is FCR, and  $\Lambda(\Omega') = \bigcup Y_i$ ,

$$I_{\Omega'} = \bigcap_{\mu \in \Lambda(\Omega')} I_\mu = \bigcap_{i=1}^r \bigcap_{\mu \in Y_i} I_\mu.$$

Because  $I_{\Omega'}$  is prime, we have that  $I_{\Omega'} = \bigcap_{\mu \in Y_i} I_\mu$ , for some  $i$ . Let  $\Omega$  denote the ideal of elements vanishing on  $X_i$ . Note that  $Y_i$  is dense in  $X_i$  for each  $i$  since  $\Lambda(\Omega')$  is dense in  $X$  (see Lemma 3.4). Therefore  $I_{\Omega'} = I_\Omega$  by [Soe90] Proposition 1. This implies that  $\theta(I_{\Omega'}) = \theta(I_\Omega)$ , and therefore by [Soe90] Theorem 1 part (i) we have  $\Omega' = w \cdot \Omega$  for some  $w \in W$ .

Because  $\Lambda(\Omega) \cap X_i$  is dense in  $X_i$ , and  $\Lambda(\Omega) \subseteq P^+$ , it follows that  $P^+ \cap \Lambda(\Omega)$  is also dense in  $X_i$ . Therefore  $\Omega$  is dominant. Since  $I_\Omega = I_{w \cdot \Omega}$ , the proof of [Soe90] Theorem 1 part (ii) shows that there exists  $w_1 \in W$  such that  $w \cdot \Omega = w_1 \cdot \Omega$  and  $I_\lambda = I_{w_1 \cdot \lambda}$ . It follows from Lemma 4.8 that  $w_1^{-1} w(B_\lambda) = B_\lambda$ . Hence to prove the result it is enough to show that  $w_1 \in W^\lambda$ .

Write  $w_1 = uv$  with  $u \in W^\lambda$ ,  $v \in W_\lambda$ . By Lemma 3.2 we have  $I_{w_1 \cdot \lambda} = I_{v \cdot \lambda}$ . Hence  $I_\lambda = I_{v \cdot \lambda}$ . By Theorem 3.3  $\tau_\Lambda(v) = \tau_\Lambda(1)$ , and this implies that  $v = 1$ , so  $w_1 = u \in W^\lambda$ .  $\square$

**Lemma 4.10.** *Suppose  $\Omega, \Omega_1 \in \text{Spec } \mathcal{S}$  and let  $\lambda, \lambda_1$  respectively be the tautological highest weights.*

- (1) *If  $\Omega_1 = v \cdot \Omega$  with  $v \in W^\lambda$  then  $B_{\lambda_1} = v B_\lambda$ .*
- (2) *If in addition,  $u \in W^{\lambda_1}$  then  $uv \in W^\lambda$ .*

*Proof.* (1) From remark 3.1 we see that

$$v(R_\lambda) = \{\alpha \in R \mid v^{-1} H_\alpha - m \in \Omega \text{ for some } m \in \mathbb{Z}\}$$

and one can check that  $v(R_\lambda) = R_{\lambda_1}$ . Since  $v \in W^\lambda$ ,

$$v B_\lambda \subseteq R^+ \cap R_{\lambda_1} = R_{\lambda_1}^+.$$

This fact implies that  $v R_\lambda^+ \subseteq R_{\lambda_1}^+$ , and therefore  $R_{\lambda_1} \cap \mathbb{N} v B_\lambda = R_{\lambda_1}^+$ . Since there is a unique choice of simple roots in  $R_{\lambda_1}^+$ , the result follows.

(2) If  $\alpha \in B_\lambda$ , then  $v\alpha \in B_{\lambda_1}$  by part (1). Hence  $uv\alpha \in R^+$  for all such  $\alpha$ , so  $uv \in W^\lambda$ .  $\square$

*Proof of the Main Theorem.* (1) this follows from Theorem 4.9.

For (2) assume  $\Omega$  is dominant, and  $w \in W$ .

(b)  $\implies$  (a): If  $I_{w \cdot \Omega} = I_\Omega$ , then  $U_{w \cdot \Omega}$  is FCR by Proposition 4.3.

(c)  $\implies$  (b): This follows from Lemma 4.4.

(a)  $\implies$  (c): Suppose that  $\mathcal{U}_{w \cdot \Omega}$  is FCR. By Theorem 4.9, there is a dominant prime  $\Omega_1$ , and  $u \in W^{\lambda_1}$  such that  $w \cdot \Omega = u \cdot \Omega_1$ , where  $\lambda_1$  is the tautological highest weight corresponding to  $\Omega_1$ . If  $v = u^{-1} w$ , then  $\Omega$  and  $v \cdot \Omega$  are dominant, so by Theorem 4.6,  $v \in W^\lambda$ . By Lemma 4.10 we have  $w = uv \in W^\lambda$ .  $\square$

## 5. INVARIANT DIFFERENTIAL OPERATORS

An interesting class of examples arises from dual pairs ([GW98] Section 4.5, [How89], [LS89]). The application of our method is fairly routine, so we give only the most interesting examples rather than conduct an exhaustive study.

Let  $M_{p,k}$  denote the set of  $p \times k$  matrices over  $F$ , and consider cases  $A, B, C$  as follows.

| Case | $\mathbf{K}$ | $\mathbf{V}$             | Action of $K$ on $\mathbf{V}$       | $\mathfrak{g}$        |
|------|--------------|--------------------------|-------------------------------------|-----------------------|
| $A$  | $GL_k(F)$    | $M_{p,k} \times M_{k,q}$ | $(g, (a, b)) \mapsto (ag^{-1}, gb)$ | $\mathfrak{gl}_{p+q}$ |
| $B$  | $O_k(F)$     | $M_{k,n}$                | $(g, a) \mapsto ga$                 | $\mathfrak{sp}_{2n}$  |
| $C$  | $Sp_{2k}(F)$ | $M_{k,n}$                | $(g, a) \mapsto ga$                 | $\mathfrak{so}_{2n}$  |

In case  $A$  we set  $n = p + q$ . If  $p = 0$  (resp.  $q = 0$ ) we set  $V = M_{k,n}$  (resp.  $V = M_{n,k}$ ). In all cases the algebra of invariant differential operators  $\mathcal{D}(V)^K$  is generated (as an associative algebra) by a Lie algebra isomorphic to  $\mathfrak{g}$ . Hence there is a surjective homomorphism

$$\phi : U(\mathfrak{g}) \longrightarrow \mathcal{D}(V)^K. \quad (4)$$

The image of  $\mathfrak{g}$  under  $\phi$  is described explicitly on pages 69–70 of [LS89].

Let  $T$  be a maximal torus of  $K$ . In [MR], it is shown that if zero is not a  $T$ -weight of  $V$  then  $\mathcal{D}(V)^K$  has enough finite dimensional representations. Thus,  $\mathcal{D}(V)^K$  is FCR since it is an image of  $\mathcal{U}(\mathfrak{g})$ . We remark that zero is not a weight of  $V$  in cases  $A$  and  $C$  of the above table and in case  $B$  for even  $k$ .

There is a multiplicity free decomposition of  $\mathcal{O}(V)$  as a  $F[K] \otimes \mathcal{U}(\mathfrak{g})$ -module ([GW98] Theorem 4.5.14). Furthermore, as a  $\mathcal{U}(\mathfrak{g})$ -module,  $\mathcal{O}(V)$  is a direct sum of simple highest weight modules. Let

$$\Lambda = \{\lambda \in \mathfrak{h}^* \mid \mathcal{O}(V) \text{ has a } \mathfrak{g}\text{-submodule isomorphic to } L(\lambda)\}.$$

If  $\Omega$  is a radical ideal of  $\mathcal{S}$ , and  $\Omega_1, \dots, \Omega_t$  are the prime ideals of  $\mathcal{S}$  that are minimal over  $\Omega$ , we set  $I_\Omega = \bigcap_{i=1}^t I_{\Omega_i}$ .

**Lemma 5.1.** *If  $\Omega$  is the radical ideal of  $\mathcal{S}$  such that  $\mathcal{V}(\Omega) = \overline{\Lambda}$ , the Zariski closure of  $\Lambda$ , then*

$$\ker \phi = I_\Omega. \quad (5)$$

*Proof.* Suppose that  $\overline{\Lambda} = X_1 \cup \dots \cup X_t$  is the decomposition of  $\overline{\Lambda}$  into irreducible components, and for  $1 \leq i \leq t$ , let  $\Omega_i$  be the prime ideals of  $\mathcal{S}$  such that  $\mathcal{V}(\Omega_i) = X_i$ . By Lemma 3.4,  $\Lambda_i = \Lambda \cap X_i$  is dense in  $X_i$ , and so by ([Soe90] Proposition 1) we have:

$$I_{\Omega_i} = \bigcap_{\lambda \in \Lambda_i} \text{ann}_{\mathcal{U}} L(\lambda)$$

for  $1 \leq i \leq t$ . Therefore, since  $\mathcal{O}(V)$  is a faithful  $\mathcal{D}(V)^K$ -module and  $\Lambda = \bigcup_{i=1}^t \Lambda_i$  we have:

$$\begin{aligned} \ker \phi &= \{u \in \mathcal{U} \mid \phi(u)\mathcal{O}(V) = 0\} \\ &= \bigcap_{\lambda \in \Lambda} \text{ann}_{\mathcal{U}} L(\lambda) \\ &= \bigcap_{i=1}^t \bigcap_{\lambda \in \Lambda_i} \text{ann}_{\mathcal{U}} L(\lambda) \\ &= I_{\Omega}. \end{aligned}$$

□

Note that  $I_{\Omega}$  is a completely prime ideal. We identify a situation where it is zero.

**Theorem 5.2.** *Assume that  $\text{rank } \mathfrak{g} \leq \text{rank } K$  then  $\mathcal{U} \cong \mathcal{D}(V)^K$ .*

*Proof.* Let  $\{\mathcal{U}_N\}$  be the usual filtration on  $\mathcal{U}$  and  $\{D_N\}$  the Bernstein filtration on  $\mathcal{D}(V)$ . Then  $K$  preserves each  $D_N$ , so acts on  $\text{gr}\mathcal{D}(V)$  and  $\text{gr}(\mathcal{D}(V)^K) = (\text{gr}\mathcal{D}(V))^K$ . Now  $\mathfrak{g}$  maps onto  $D_2^K$  and this induces a surjection  $\mathcal{U}_N \rightarrow D_{2N}^K$  with kernel  $I_{\Omega} \cap \mathcal{U}_N$ . Passing to the associated graded rings we obtain a surjection

$$\mathcal{S}(\mathfrak{g}) = \text{gr}\mathcal{U} \rightarrow \text{gr}\mathcal{D}(V)^K = \mathcal{S}(V \oplus V^*)^K$$

with kernel  $\text{gr}I_{\Omega}$ . Then if  $\text{rank } K \geq \text{rank } \mathfrak{g}$  then we can apply the Second Fundamental Theorem of Invariant Theory in the free case ([GW98] Corollary 4.2.5) to  $V \oplus V^*$ . We conclude that  $\text{gr}I_{\Omega} = (0)$  so  $I_{\Omega} = (0)$ . □

Our next aim is to describe the irreducible constituents of  $\mathcal{O}(V)$  as a  $K$ -module and as a  $\mathcal{U}(\mathfrak{g})$ -module. With this goal in mind we set up some standard notation.

By a partition we will mean a finite sequence of weakly decreasing non-negative integers. We will denote partitions by lower case Greek letters. The *length* of a partition is denoted  $\ell(\alpha) = \max\{i \mid \alpha_i > 0\}$ . The *conjugate* of a partition  $\lambda$  is a partition  $\lambda'$  whose  $i^{\text{th}}$  part is given by  $|\{j \mid \lambda_j \geq i\}|$ . Note that  $\ell(\lambda) = \lambda'_1$ .

The highest weights of representations of  $\mathfrak{g}$  (resp.  $K$ ) are given by  $m$ -tuples where  $m$  is the rank of  $\mathfrak{g}$  (resp.  $K$ ). The  $m$ -tuple  $(a_1, \dots, a_m)$  corresponds to the weight  $\sum_{i=1}^m a_i \epsilon_i$  where the  $\epsilon_i$  are defined in Section 2.3.1 of [GW98].

At this point we consider the cases separately.

**5.1. Case A.** As a basis for  $\mathfrak{h}$  we take  $\{E_1, \dots, E_n\}$  where  $E_i$  is the  $n \times n$  matrix with a 1 in the row  $i$ , column  $i$  entry and zeros elsewhere.

Given non-negative integer partitions  $\alpha$  and  $\beta$  with  $\ell(\alpha) \leq p$ ,  $\ell(\beta) \leq q$ , and  $\ell(\alpha) + \ell(\beta) \leq k$ , let  $V_{(\alpha, \beta)}$  denote the irreducible  $\mathfrak{gl}_n(F)$  module with highest weight:

$$(\alpha, \beta)_{\mathfrak{g}} := \underbrace{(-k - \alpha_p, -k - \alpha_{p-1}, \dots, -k - \alpha_1, \beta_1, \beta_2, \dots, \beta_q)}_n.$$

Let  $V^{(\alpha, \beta)}$  denote the irreducible representation of  $GL_k$  with highest weight:

$$(\alpha, \beta)_K := \underbrace{(\alpha_1, \alpha_2, \dots, \alpha_p, 0, \dots, 0, -\beta_q, \dots, -\beta_2, -\beta_1)}_k.$$

**Lemma 5.3.** *There is a multiplicity free decomposition under the joint action of  $K$  and  $\mathfrak{g}$  we have the multiplicity free decomposition:*

$$\mathcal{O}(V) = \bigoplus V^{(\alpha, \beta)} \otimes V_{(\alpha, \beta)},$$

where the direct sum is over the set

$$W_k(p, q) := \left\{ (\alpha, \beta) \left| \begin{array}{l} \alpha \text{ and } \beta \text{ partitions such that} \\ \ell(\alpha) \leq p, \ell(\beta) \leq q, \quad \text{and} \\ \ell(\alpha) + \ell(\beta) \leq k \end{array} \right. \right\}$$

*Proof.* See [EW04] p. 356.  $\square$

Define  $\Lambda_k(p, q) = \{(\alpha, \beta)_{\mathfrak{g}} | (\alpha, \beta) \in W_k(p, q)\}$ , and let  $Z_k(p, q) := \overline{\Lambda_k(p, q)}$  denote the Zariski closure of these weights.

If  $k < n$  define:

$$\Omega_m := \begin{cases} (E_{k+1}, E_{k+2}, \dots, E_n), & \text{if } m = 0; \\ (E_1 + k, \dots, E_m + k, E_{m+k+1}, \dots, E_n), & \text{if } 0 < m < n - k; \\ (E_1 + k, \dots, E_{n-k} + k), & \text{if } m = n - k. \end{cases}$$

**Lemma 5.4.** *The decomposition of  $Z_k(p, q)$  into irreducible components is as follows:*

- (1)  $Z_k(p, q) = \mathfrak{h}^*$ , if  $n \leq k$ .
- (2) If  $k < n \leq 2k$ , then

$$Z_k(p, q) = \bigcup_{m \in \Phi_p} \mathcal{V}(\Omega_m)$$

where:

$$\Phi_p = \begin{cases} \{0, \dots, p\}, & \text{if } p \leq n - k; \\ \{0, \dots, n - k\}, & \text{if } n - k \leq p \leq k; \\ \{p - k, \dots, n - k\}, & \text{if } k \leq p \leq n. \end{cases}$$

- (3) If  $2k \leq n$  then

$$Z_k(p, q) = \bigcup_{m \in \Phi_p} \mathcal{V}(\Omega_m)$$

where:

$$\Phi_p = \begin{cases} \{0, \dots, p\}, & \text{if } p \leq k; \\ \{p - k, \dots, p\}, & \text{if } k \leq p \leq n - k; \\ \{p - k, \dots, n - k\}, & \text{if } p \geq n - k. \end{cases}$$

*Proof.* Statement (1) follows because if  $n \leq k$ , then  $\mathcal{O}(V)$  contains the irreducible  $\mathfrak{gl}_n(F)$  module with highest weight  $(\alpha, \beta)_{\mathfrak{g}}$  for any partitions  $\alpha, \beta$  with  $\ell(\alpha) \leq p$ , and  $\ell(\beta) \leq q$ . For statements (2) and (3) we define, for  $0 \leq i \leq p$

$$W_k^i(p, q) = \{(\alpha, \beta) \in W_k(p, q) | \ell(\alpha) = p - i\},$$

and

$$\Lambda_k^i(p, q) = \{(\alpha, \beta)_{\mathfrak{g}} | (\alpha, \beta) \in W_k(p, q)\}.$$

Suppose  $(\alpha, \beta) \in W_k^i(p, q)$ . Then,  $0 \leq i \leq p$  since  $0 \leq \ell(\alpha) \leq p$ . Also,  $\ell(\alpha) \leq k$  implies that  $p - k \leq i$ . In addition, the set of weights  $(\alpha, \beta)_{\mathfrak{g}}$  such that  $\ell(\alpha) = p - i$  and  $\ell(\beta) < k - \ell(\alpha) = k + i - p$  is contained in the Zariski closure of the set of weights  $(\alpha, \beta)_{\mathfrak{g}}$  with  $\ell(\beta) = k + i - p$ . Since we require that  $\ell(\beta) \leq q$ , we can restrict our attention to the highest weights  $(\alpha, \beta)_{\mathfrak{g}}$  such that  $\ell(\alpha) = p - i$  and  $k + i - p \leq q$ .

That is,  $i \leq n - k$ . It follows that  $Z_k(p, q) = \bigcup \overline{\Lambda_k^i(p, q)}$  where the union is over all  $i$  such that  $\max(0, p - k) \leq i \leq \min(p, n - k)$ . It is easy to check that for these values of  $i$ ,  $\overline{\Lambda_k^i(p, q)} = \mathcal{V}(\Omega_i)$ . The result follows by considering individual cases.  $\square$

**Proposition 5.5.** *If  $n$  and  $k$  are fixed, then for  $0 \leq i < j \leq n - k$  we have  $I_{\Omega_i} = I_{\Omega_j}$ . Moreover,  $\Omega_0$  and  $\Omega_{n-k}$  are dominant.*

*Proof.* To see that  $\Omega_0$  is dominant note that

$$\sum_{i=1}^k \mathbb{N}\bar{\omega}_i \subseteq \mathcal{V}(\Omega_0) = \sum_{i=1}^k \mathbb{F}\bar{\omega}_i.$$

Similarly  $\Omega_{n-k}$  is dominant. Now set  $\Omega = \Omega_{n-k}$ , so that:

$$\Omega = (E_1 + k, E_2 + k, \dots, E_{n-k} + k),$$

and let  $\lambda = \lambda_\Omega$ . For  $0 \leq i \leq n - k$  we define  $w_i \in W = S_n$  by:

$$w_i(j) = \begin{cases} j, & \text{for } 1 \leq j \leq i; \\ j + k, & \text{for } i + 1 \leq j \leq n - k; \\ j - n + k + i, & \text{for } n - k + 1 \leq j \leq n. \end{cases}$$

Then, it is easy to see that  $w_i.\Omega = \Omega_i$ , and that  $w_i \in W^\lambda$ . The result follows from the Lemma 4.4.  $\square$

*Remark 5.6.* Explicit generators for  $I_{\Omega_0}$  are given in [Pro04] using Capelli identities.

**5.2. Cases B and C.** We first turn our attention to Case B.

**Lemma 5.7.** *There is a multiplicity free decomposition under the joint action of  $K$  and  $\mathfrak{g}$  given by*

$$\mathcal{O}(V) = \bigoplus_{\mu \in M} V^\mu \otimes V_\mu,$$

where we index the summands by the partitions  $\mu$  in the set:

$$M := \{\mu \text{ is a partition such that } \mu'_1 + \mu'_2 \leq k \text{ and } \ell := \mu'_1 \leq n\}.$$

The highest weight of the  $sp_{2n}$ -module,  $V_\mu$  is given by:

$$\underbrace{\left(-\frac{k}{2}, -\frac{k}{2}, \dots, -\frac{k}{2} - \mu_\ell, \dots, -\frac{k}{2} - \mu_1\right)}_n$$

*Proof.* See [EW04] p. 353.  $\square$

We describe the Zariski closure,  $Z$  in  $\mathfrak{h}^*$  of the set of weights of  $\mathcal{O}(V)$  as a  $\mathfrak{g}$ -module. If  $k \geq 2n$  then  $Z = \mathfrak{h}^*$ , so we consider the case where  $k < 2n$ . For  $k$  odd, we can see using Theorem 4.1 that  $\mathcal{D}(V)^K$  has no finite dimensional modules, so we consider the  $k$  even case.

Suppose that  $k = 2p$ . Fix  $i$  with  $1 \leq i \leq \min(p, n - p)$  and set  $r = n - p - i$ ,  $s = n - p + i$ . Then define

$$\Omega_i = (H_1, H_2, \dots, H_{r-1}, H_r + 1, H_{r+1}, \dots, H_{s-1}, H_s + H_{s+1} + \dots + H_n + p + 1).$$

We also set

$$\Omega_0 = (H_1, \dots, H_{n-p-1}, H_{n-p} + H_{n-p+1} + \dots + H_n + p).$$

**Lemma 5.8.** *The decomposition of  $Z$  into irreducible components is given by*

$$Z = \bigcup_{i=0}^{\min(p, n-p)} \mathcal{V}(\Omega_i).$$

*Proof.* For  $0 \leq i \leq \min(p, n-p)$ , set

$$M_i = \{\mu \text{ is a partition such that } \mu'_1 + \mu'_2 \leq k, \text{ and } \mu'_1 = p + i\}.$$

and let  $\Lambda_i \subseteq \mathfrak{h}^*$  be the set of highest weights of the modules  $V_\mu$  with  $\mu \in M_i$ . Note that if  $\mu'_1 + \mu'_2 \leq k = 2p$  and  $\mu'_1 < p$  then the highest weight of  $V_\mu$  is in  $\bar{\Lambda}_0$ .

If  $0 \leq i < j \leq \min(p, n-p)$  neither of the ideals  $\Omega_i, \Omega_j$  is contained in the other. Therefore, it suffices to show that the Zariski closure of  $\Lambda_i$  is  $\mathcal{V}(\Omega_i)$ . Suppose that  $\mu \in M_i$  and let  $\alpha = \sum_{i=1}^n a_i \epsilon_i$  be the highest weight of  $V_\mu$ . Then  $a_i = -p$  for  $1 \leq i \leq r = n - p - i$ , and  $a_i = -p - 1$  for  $r + 1 \leq i \leq s = n - p + i$ . The result follows easily.  $\square$

Note that  $\text{GK-dim } \mathcal{S}/\Omega_i = p - i$ , where  $\text{GK-dim}$  is the Gelfand-Kirillov dimension. Let  $I_i = I_{\Omega_i}$  for  $0 \leq i \leq p$ .

Recall the homomorphism  $\phi$  from equation (4).

**Lemma 5.9.** *We have  $\ker \phi = I_0$ .*

*Proof.* By Lemma 5.1,  $\ker \phi = \cap_{i=0}^p I_i$ . Since  $\ker \phi$  is prime, it follows that  $I_i \subseteq \ker \phi \subseteq I_0$  for some  $i$ . Hence  $I_i \cap \mathcal{Z} \subseteq I_0 \cap \mathcal{Z}$ , that is  $\Omega_i \cap \mathcal{Z} \subseteq \Omega_0 \cap \mathcal{Z}$ . If  $i \geq 1$  then since  $\mathcal{S}$  is a finitely generated  $\mathcal{Z}$ -module we would have:

$$\begin{aligned} p - i &= \text{GK-dim}(\mathcal{S}/\Omega_i) = \text{GK-dim}(\mathcal{Z}/\mathcal{Z} \cap \Omega_i) \\ &\geq \text{GK-dim}(\mathcal{Z}/\mathcal{Z} \cap \Omega_0) = p \end{aligned}$$

a contradiction.  $\square$

We show next that  $\mathcal{U}/I_0$  is FCR. To do this define

$$\Omega = (H_{p+1}, \dots, H_n),$$

and define  $w \in W$  so that if  $a = (a_1, \dots, a_n) \in \mathfrak{h}^*$ , we have  $w(a) = b$ , where:

$$b_i = \begin{cases} a_{i+p}, & \text{for } 1 \leq i \leq n-p; \\ a_{i+p-n}, & \text{for } n-p+1 \leq i \leq n. \end{cases}$$

It is easy to check that  $\Omega$  is dominant,  $w.\Omega = \Omega_0$  and  $w \in W^\lambda$  where  $\lambda = \lambda_\Omega$ . Thus by the main theorem,  $I_0 = I_\Omega$  and  $\mathcal{U}_\Omega$  is FCR. We expect that

$$I_0 \subset I_1 \subset \dots \subset I_p$$

with all inclusions strict. If  $p = 1$  this follows from Lemmas 5.1 and 5.9.

Case C is relatively uninteresting, the Zariski closure  $\mathcal{V}(\Omega)$  of the set of highest weight occurring in  $\mathcal{O}(V)$  as a  $\mathfrak{g}$ -module is irreducible. Also  $\mathcal{U}_\Omega \cong D(V)^K$  has enough finite dimensional modules from [MR] and thus is FCR. Furthermore,  $\mathcal{V}(\Omega) = \mathfrak{h}^*$  if and only if  $n \leq k$ .

## REFERENCES

- [BJ01] Walter Borho and Anthony Joseph. Sheets and topology of primitive spectra for semisimple Lie algebras *J. Algebra*, 259(1):310–311, 2003.; Corrigendum to *J. Algebra* **244** (2001), no. 1, 76–167.
- [Dix70] J. Dixmier. Idéaux primitifs dans l’algèbre enveloppante d’une algèbre de Lie semi-simple complexe. *C. R. Acad. Sci. Paris Sér. A-B*, 271:A134–A136, 1970.
- [EW04] T. J. Enright and J. F. Willenbring. Hilbert series, Howe duality and branching for classical groups. *Ann. of Math. (2)*, 159(1):337–375, 2004.
- [GW98] R. Goodman and N.R. Wallach. *Representations and invariants of the classical groups*. Cambridge University Press, Cambridge, 1998.
- [HC] Harish-Chandra, On representations of Lie algebras, *Ann. of Math. (2)*, 50, 1949, 900–915.
- [How89] R. Howe. Remarks on classical invariant theory. *Trans. Amer. Math. Soc.*, 313(2):539–570, 1989.
- [Hu90] J.E. Humphreys, *Reflection groups and Coxeter groups*. Cambridge Univ. Press, Cambridge, 1990.
- [Jan83] J. C. Jantzen. *Einhüllende Algebren halbeinfacher Lie-Algebren*, volume 3 of *Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)]*. Springer-Verlag, Berlin, 1983.
- [Jo98] A. Joseph. Sur l’annulateur d’un module de Verma, pages 237–300, *Representation theories and algebraic geometry (Montreal, PQ, 1997)*, NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci., Kluwer Acad. Publ., Dordrecht, 1998.
- [KS94] H. Kraft and L. W. Small. Invariant algebras and completely reducible representations. *Math. Res. Lett.*, 1(3):297–307, 1994.
- [LS89] T. Levasseur and J. T. Stafford. Rings of differential operators on classical rings of invariants. *Mem. Amer. Math. Soc.*, 81(412):vi+117, 1989.
- [MR] I. M. Musson and S. L. Rueda. Finite dimensional representations of invariant differential operators. *Trans. Amer. Math. Soc.* 357 (2004), no. 7, 2739–2752.
- [P05] Nikolaos Papalexiou. Order relations on the induced and prime ideals in the enveloping algebra of a semisimple Lie algebra *Journal of Algebra* 302 (2006) 1–15.
- [P] Nikolaos Papalexiou. On the prime spectrum of the enveloping algebra and characteristic varieties. To Appear in the *Journal of Algebra and its Applications*.
- [Pro04] V. Protsak. Rank ideals and Capelli identities. *J. Algebra* 273 (2004), no. 2, 686–699.
- [Soe90] W. Soergel. The prime spectrum of the enveloping algebra of a reductive Lie algebra. *Math. Z.*, 204(4):559–581, 1990.
- [Soe] W. Soergel.  
<http://home.mathematik.uni-freiburg.de/soergel/PReprints/Korrektur.pdf>

IAN M. MUSSON, UNIVERSITY OF WISCONSIN - MILWAUKEE, DEPARTMENT OF MATHEMATICAL SCIENCES, P.O. BOX 0413, MILWAUKEE WI 53201-0413  
*E-mail address:* `musson@uwm.edu`

JEB F. WILLENBRING, UNIVERSITY OF WISCONSIN - MILWAUKEE, DEPARTMENT OF MATHEMATICAL SCIENCES, P.O. BOX 0413, MILWAUKEE WI 53201-0413  
*E-mail address:* `jw@uwm.edu`